



Riverdale®
RAZOR®
HERBICIDE

**THE COMPLETE BROAD
SPECTRUM POSTEMERGENCE
PROFESSIONAL HERBICIDE FOR
INDUSTRIAL, FORESTRY, TURF, VEGETATION
MANAGEMENT AND ORNAMENTAL WEED CONTROL**

ACTIVE INGREDIENT:

Glyphosate, N-(phosphonomethyl) glycine,
in the form of its isopropylamine salt* 41.0%

OTHER INGREDIENTS: 59.0%

TOTAL 100.0%

* Contains 480 grams per litre or 4 pounds per U.S. gallon of the active ingredient glyphosate, in the form of its isopropylamine salt. Equivalent to 356 grams per litre or 3 pounds per U.S. gallon of the acid, glyphosate.

**KEEP OUT OF REACH OF CHILDREN
CAUTION - CAUCION**

Si usted no entiende la etiqueta, busque a alguien para que se la explique a usted en detalle.
(If you do not understand the label, find someone to explain it to you in detail.)

SEE INSIDE BOOKLET FOR FIRST AID AND ADDITIONAL PRECAUTIONARY STATEMENTS

**For Chemical Spill, Leak, Fire, or Exposure, Call CHEMTREC (800) 424-9300
For Medical Emergencies Only, Call (877) 325-1840**

**MANUFACTURED BY
NUFARM AMERICAS INC.,
BURR RIDGE, IL 60527-0866**

**EPA REG. NO. 228-366
EPA EST. NO. 228-IL-1**



**PRECAUTIONARY STATEMENTS
HAZARDS TO HUMANS AND DOMESTIC ANIMALS
CAUTION - CAUCION**

Causes moderate eye irritation. Harmful if swallowed or inhaled. Do not get in eyes or on clothing. Avoid breathing vapor or spray mist.

PERSONAL PROTECTIVE EQUIPMENT (PPE)

For uses covered under the Worker Protection Standard (WPS), 40 CFR Part 170 - in general, only agricultural plant uses are covered by the WPS: applicators and other handlers must wear long-sleeved shirt and long pants, shoes plus socks, and protective eyewear. Discard clothing and other absorbent materials that have been drenched or heavily contaminated with this product's concentrate. Do not reuse them. Follow manufacturer's instructions for cleaning/maintaining PPE. If no such instructions for washables, use detergent and hot water. Keep and wash PPE separately from other laundry.

USER SAFETY RECOMMENDATIONS

USERS SHOULD:

- Wash hands before eating, drinking, chewing gum, using tobacco or using the toilet.
- Remove clothing immediately if pesticide gets inside. Then wash thoroughly and put on clean clothing.
- Remove PPE immediately after handling this product. Wash the outside of gloves before removing. As soon as possible, wash thoroughly and change into clean clothing.

FIRST AID

IF IN EYES	<ul style="list-style-type: none"> • Hold eye open and rinse slowly and gently with water for 15 to 20 minutes. • Remove contact lenses, if present, after the first 5 minutes, then continue rinsing eye. • Call a poison control center or doctor for treatment advice.
IF SWALLOWED	<ul style="list-style-type: none"> • Call a poison control center or doctor immediately for treatment advice. • Have person sip a glass of water if able to swallow. • Do not induce vomiting unless told to do so by a poison control center or doctor. Do not give anything by mouth to an unconscious person.
IF INHALED	<ul style="list-style-type: none"> • Move person to fresh air. • If person is not breathing, call 911 or an ambulance then give artificial respiration, preferably mouth-to-mouth if possible. • Call a poison control center or doctor for further treatment advice.

HOT LINE NUMBER

Have the product container or label with you when calling a poison control center or doctor, or going for treatment. You may also contact 1-877-325-1840 for emergency medical treatment information.

ENGINEERING CONTROL STATEMENTS

When handlers use closed systems, enclosed cabs, or aircraft in a manner that meets the requirements listed in Worker Protection Standard (WPS) for agricultural pesticides [40 CFR 170.240 (d) (4-6)], the handler PPE requirements may be reduced or modified as specified in the WPS.

ENVIRONMENTAL HAZARDS

Do not apply directly to water, to areas where surface water is present or to intertidal areas below the mean high water mark. Do not contaminate water when disposing of equipment washwaters.

PHYSICAL OR CHEMICAL HAZARDS

Spray solutions of this product should be mixed, stored and applied using only stainless steel, aluminum, fiberglass, plastic or plastic-lined steel containers.

DO NOT MIX, STORE OR APPLY THIS PRODUCT OR SPRAY SOLUTIONS OF THIS PRODUCT IN GALVANIZED STEEL OR UNLINED STEEL (EXCEPT STAINLESS STEEL) CONTAINERS OR SPRAY TANKS. This product or spray solutions of this product react with such containers and tanks to produce hydrogen gas which may form a highly combustible gas mixture. This gas mixture could flash or explode, causing serious personal injury, if ignited by open flame, spark, welder's torch, lighted cigarette or other ignition source.

DIRECTIONS FOR USE

It is a violation of Federal law to use this product in a manner inconsistent with its labeling. **READ ENTIRE LABEL BEFORE USING THIS PRODUCT.** Do not apply this product in a way that will contact workers or other persons, either directly or through drift. Only protected handlers may be in the area during application. For any requirements specific to your State or Tribe, consult the agency responsible for pesticide regulations.

AGRICULTURAL USE REQUIREMENTS

Use this product only in accordance with its labeling and with the Worker Protection Standard, 40 CFR part 170. This Standard contains requirements for the protection of agricultural workers on farms, forests, nurseries, and greenhouses, and handlers of agricultural pesticides. It contains requirements for training, decontamination, notification, and emergency assistance. It also contains specific instructions and exceptions pertaining to the statements on this label about Personal Protective Equipment (PPE) and restricted-entry interval. The requirements in this box only apply to uses of this product that are covered by the Worker Protection Standard.

Do not enter or allow worker entry into treated areas during the restricted entry interval (REI) of 4 hours.

PPE required for early entry to treated areas that is permitted under the Worker Protection Standard and that involves contact with anything that has been treated, such as plants, soil, or water, is: Coveralls, chemical-resistant gloves made of any waterproof material such as barrier laminate or vitron >13, shoes plus socks, and protective eyewear.

NON-AGRICULTURAL USE REQUIREMENTS

The requirements in this box apply to uses of this product that are NOT within the scope of the Worker Protection Standard for agricultural pesticides (40 CFR part 170). The WPS applies when this product is used to produce agricultural plants on farms, forests, nurseries, or greenhouses.

Keep people and pets off treated areas until spray solution has dried.

GENERAL INFORMATION

Do not apply this product using aerial spray equipment except under conditions as specified within this label.

This product, a water-soluble liquid, mixes readily with water to be applied as a foliar spray for the control or elimination of most herbaceous plants. It may be applied through most standard industrial or field-type sprayers after dilution and thorough mixing with water in accordance with label instructions.

This product moves through the plant from the point of foliage contact to and into the root system. Visible effects on most annual weeds occur within 2 to 4 days, but on most perennial weeds may not occur for 7 days or more. Extremely cool or cloudy weather following treatment may slow activity of this product and delay visual effects of control. Visible effects are a gradual wilting and yellowing of the plant which advances to complete browning of above-ground growth and deterioration of underground plant parts.

Unless otherwise specified on this label, delay application until vegetation has emerged and reached the stages described for control of such vegetation under the "WEEDS CONTROLLED" section of this label.

Always use the higher rate of this product per acre within the recommended range when (1) weed growth is heavy or dense, or (2) weeds are growing in an undisturbed (non-cultivated) area.

Do not treat weeds under poor growing conditions such as drought stress, disease or insect damage, as reduced weed control may result. Reduced results may also occur when treating weeds heavily covered with dust.

Reduced control may result when applications are made to annual or perennial weeds that have been mowed, grazed, or cut, and have not been allowed to regrow to the recommended stage for treatment.

Rainfall or irrigation occurring within 6 hours after application may reduce effectiveness. Heavy rainfall or irrigation within 2 hours after application may wash the spray mixture off the foliage and a repeat treatment may be required.

This product does not provide residual weed control. For subsequent residual weed control, follow a label-approved herbicide program. Read and carefully observe the cautionary statements and all other information appearing on the labels of all herbicides used.

For best results, spray coverage should be uniform and complete. Do not spray weed foliage to the point of runoff.

DOMESTIC ANIMALS: This product is considered to be relatively nontoxic to dogs and other domestic animals; however, ingestion of this product or large amounts of freshly sprayed vegetation may result in temporary gastrointestinal irritation (vomiting, diarrhea, colic, etc.). If such symptoms are observed, provide the animal with plenty of fluids to prevent dehydration. Call a veterinarian if symptoms persist for more than 24 hours.

If application rates for grass seed, sod production, general noncrop areas, industrial sites, pasture grass and rangeland total 3 quarts per acre or less, no waiting period between treatment and feeding or livestock grazing is required. If the rate is greater than 3 quarts per acre, remove domestic livestock before application and wait 8 weeks after application before grazing or harvesting.

ATTENTION

AVOID DRIFT. EXTREME CARE MUST BE USED WHEN APPLYING THIS PRODUCT TO PREVENT INJURY TO DESIRABLE PLANTS AND CROPS.

Do not allow the herbicide solution to mist, drip, drift or splash onto desirable vegetation since minute quantities of this product can cause severe damage or destruction. The likelihood of injury occurring from the use of this product is greatest when winds are gusty or in excess of 5 miles per hour or when other conditions, including lesser wind velocities, will allow spray drift to occur. When spraying, avoid combinations of pressure and nozzle type that will result in splatter or fine particles (mist) which are likely to drift. **AVOID SPRAYING AT EXCESSIVE SPEED OR PRESSURE.**

NOTE: Use of this product in any manner not consistent with this label may result in injury to persons, animals or crops, or other unintended consequences. Keep container closed to prevent spills and contamination.

For questions or additional rate information contact your Nufarm representative.

MIXING, ADDITIVES AND APPLICATION INSTRUCTIONS

APPLY THESE SPRAY SOLUTIONS IN PROPERLY MAINTAINED AND CALIBRATED EQUIPMENT CAPABLE OF DELIVERING DESIRED VOLUMES. DO NOT APPLY WHEN WIND OR OTHER CONDITIONS FAVOR DRIFT. HAND-GUN APPLICATIONS SHOULD BE PROPERLY DIRECTED TO AVOID SPRAYING DESIRABLE PLANTS.

NOTE: REDUCED RESULTS MAY OCCUR IF WATER CONTAINING SOIL IS USED, SUCH AS WATER FROM PONDS AND UNLINED DITCHES.

MIXING

Use caution to avoid siphoning back into the carrier source. Use approved anti-back-siphoning devices where required by state or local regulations. During mixing and application, foaming of the spray solution may occur. To prevent or minimize foam, avoid the use of mechanical agitators, terminate by-pass and return lines at the bottom of the tank and, if needed, use an approved anti-foam or defoaming agent.

TANK MIXTURES

Always predetermine the compatibility of labeled tank mixtures of this product with water carrier by mixing small proportional quantities in advance.

Mix labeled tank mixtures of this product with water as follows:

Add individual formulations to the spray tank as follows: wettable powder, flowable, emulsifiable concentrate, drift control additive, water soluble liquid followed by surfactant.

Maintain good agitation at all times until the contents of the tank are sprayed. If the spray mixture is allowed to settle, thorough agitation is required to resuspend the mixture before spraying is resumed.

Carefully select proper nozzle to avoid spraying a fine mist. For best results with conventional ground application equipment, use flat fan nozzles.

Clean sprayer and parts immediately after using this product by thoroughly flushing with water.

MIXING FOR HAND-HELD SPRAYERS

Prepare the desired volume of spray solution by mixing the amount of this product in water as shown in the following table:

SPRAY SOLUTION

DESIRED VOLUME	AMOUNT OF RIVERDALE RAZOR HERBICIDE					
	1/2%	1%	1-1/2%	2%	5%	10%
1 Gal.	2/3 fluid ounce	1-1/3 fluid ounces	2 fluid ounces	2-2/3 fluid ounces	6-1/2 fluid ounces	13 fluid ounces
25 Gal.	1 pint	1 quart	1-1/2 quart	2 quarts	5 quarts	10 quarts
100 Gal.	2 quarts	1 gallon	1-1/2 gallon	2 gallons	5 gallons	10 gallons

2 tablespoons = 1 fluid ounce

For use in knapsack sprayers, it is suggested that the recommended amount of this product be mixed with water in a larger container. Fill sprayer with the mixed solution.

ADDITIVES

SURFACTANTS

Nonionic surfactants which are labeled for use with herbicides may be used. Do not reduce rates of this product when adding surfactant. When adding additional surfactants follow manufacturers rates and recommendations for use of the surfactant. Read and carefully observe surfactant cautionary statements and other information appearing on the surfactant label.

AMMONIUM SULFATE

The addition of 1 to 2 percent dry ammonium sulfate by weight or 8.5 to 17 pounds per 100 gallons of water may increase the performance of this product, and this product plus 2,4-D, Dicamba or residual herbicide tank mixtures on annual and perennial weeds. The improvement in performance may be apparent where environmental stress is a concern. If ammonium sulfate is added directly to the spray tank, add slowly with agitation. Adding too quickly may clog outlet line. Ensure that ammonium sulfate is completely dissolved in the spray tank before adding herbicides or surfactant. Thoroughly rinse the spray system with clean water after use to reduce corrosion.

NOTE: The use of ammonium sulfate as an additive does not preclude the need for additional surfactant. When using ammonium sulfate, apply this product at rates recommended on this label. Lower rates will result in reduced performance.

COLORANTS OR DYES

Agriculturally-approved colorants or marking dyes may be added to this product. Colorants or dyes used in spray solutions of this product may reduce performance, especially at lower rates or dilutions. Use colorants or dyes according to the manufacturer's recommendations.

APPLICATION EQUIPMENT AND TECHNIQUES

Do not apply this product through any type of irrigation system.

This product may be applied with the following application equipment:

Aerial - Fixed Wing and Helicopter

Broadcast Spray

Controlled Droplet Applicator (CDA) - Hand-held or boom-mounted applicators which produce a spray consisting of a narrow range of droplet sizes.

Hand-Held and High-Volume Spray Equipment - Knapsack and backpack sprayers, pump-up pressure sprayers, handguns, handwands, mistblowers*, lances and other hand-held and motorized spray equipment used to direct the spray onto weed foliage. *This product is not registered in California or Arizona for use in mistblowers.

Selective equipment - Recirculating sprayers, shielded sprayers, hooded sprayers and wiper applicators. See the appropriate part of this section for specific instructions and rates of application

AERIAL EQUIPMENT

Use the recommended rates of this herbicide in sufficient volume of water to insure thorough coverage unless otherwise specified in this label. See the "WEEDS CONTROLLED" section of this label for specific use rates. Unless otherwise specified, do not exceed 1 quart per acre. Aerial applications of this product may be made in annual cropping conventional tillage systems, fallow and reduced tillage systems.

Avoid direct application to any body of water.

AVOID DRIFT - DO NOT APPLY DURING LOW-LEVEL INVERSION CONDITIONS, WHEN WINDS ARE GUSTY OR UNDER ANY OTHER CONDITION WHICH FAVORS DRIFT. DRIFT MAY CAUSE DAMAGE TO ANY VEGETATION CONTACTED TO WHICH TREATMENT IS NOT INTENDED. TO PREVENT INJURY TO ADJACENT DESIRABLE VEGETATION, APPROPRIATE BUFFER ZONES MUST BE MAINTAINED.

Coarse sprays are less likely to drift; therefore, do not use nozzles or nozzle configurations which dispense spray as fine spray droplets. Do not angle nozzles forward into the airstream and do not increase spray volume by increasing nozzle pressure.

Drift control additives may be used. When a drift control additive is used, read and carefully observe the cautionary statements and all other information appearing on the additive label.

Ensure uniform application - To avoid streaked, uneven or overlapped application, use appropriate marking devices.

Thoroughly wash aircraft, especially landing gear, after each day of spraying to remove residues of this product accumulated during spraying or from spills. PROLONGED EXPOSURE OF THIS PRODUCT TO UNCOATED STEEL SURFACES MAY RESULT IN CORROSION AND POSSIBLE FAILURE OF THE PART. LANDING GEAR ARE MOST SUSCEPTIBLE. The maintenance of an organic coating (paint), which meets aerospace specification MIL-C-38413, may be used to prevent corrosion.

This product plus Spyder™ (Sulfometuron methyl), Diablo™ (Dicamba) or 2,4-D tank mixtures may not be applied by air in California.

SPRAY DRIFT MANAGEMENT

Avoiding spray drift at the application site is the responsibility of the applicator. The interaction of many equipment-and-weather-related factors determine the potential for spray drift. The applicator and the grower are responsible for considering all these factors when making decisions. The following drift management requirements must be followed to avoid off-target drift movement from aerial applications. These requirements do not apply to forestry applications, public health uses or to applications using dry formulations.

1. The distance of the outer most nozzles on the boom must not exceed 3/4 the length of the wingspan or rotor.
2. Nozzles must always point backward parallel with the air stream and never be pointed downwards more than 45 degrees. Where states have more stringent regulations, they should be observed.

The applicator should be familiar with and take into account the information covered in the Aerial Drift Reduction Advisory Information.

Importance of Droplet Size

The most effective way to reduce drift potential is to apply large droplets. The best drift management strategy is to apply the largest droplets that provide sufficient coverage and control. Applying larger droplets reduces drift potential, but will not prevent drift if applications are made improperly, or under unfavorable environmental conditions (see Wind, Temperature and Humidity, and Temperature Inversion section of this label).

Controlling Droplet Size

Volume-Use high flow rate nozzles to apply the highest practical spray volume. Nozzles with higher rated flows produce larger droplets.

Pressure-Use the lower spray pressures recommended for the nozzle. Higher pressure reduces droplet size and does not improve canopy protection. When higher flow rates are needed, use higher flow rate nozzles instead of increasing pressure.

Number of nozzles-Use the minimum number of nozzles that provide uniform coverage.

Nozzle Orientation-Orienting nozzles so that the spray is released backwards, parallel to the airstream will produce larger droplets than other orientations. Significant deflection from the horizontal will reduce droplet size and increase drift potential.

Nozzle Type-Use a nozzle type that is designed for the intended application. With most nozzle types, narrower spray angles produce larger droplets. Consider using low-drift nozzles. Solid stream nozzles oriented straight back produce larger droplets than other nozzle types.

Boom Length-For some use patterns, reducing the effective boom length to less than 3/4 of the wingspan or rotor length may further reduce drift without reducing swath width.

Application-Applications should not be made at a height greater than 10 feet above the top of the largest plants unless a greater height is required for aircraft safety. Making applications at the lowest height that is safe reduces exposure of droplets to evaporation and wind.

Swath Adjustment

When applications are made with a cross-wind, the swath will be displaced downwind. Therefore, on the up and downwind edges of the field, the applicator must compensate for this displacement by adjusting the path of the aircraft upwind. Swath adjustment distance should increase, with increasing drift potential (higher wind, smaller drops, etc.).

Wind

Drift potential is lowest between wind speeds of 2 to 10 mph. However, many factors, including droplet size and equipment type determine drift potential at any given speed. Application should be avoided below 2 mph due to variable wind direction and high inversion potential. Note: Local terrain can influence wind patterns. Every applicator should be familiar with local wind patterns and how they affect drift.

Temperature And Humidity

When making applications in low relative humidity, set up equipment to produce larger droplets to compensate for evaporation. Droplet evaporation is most severe when conditions are both hot and dry.

Temperature Inversions

Applications should not occur during a temperature inversion, because drift potential is high. Temperature inversions restrict vertical air mixing, which causes small suspended droplets to remain in a concentrated cloud. This cloud can move in unpredictable directions due to the light variable winds common during inversions. Temperature inversions are characterized by increasing temperatures with altitude and are common on nights with limited cloud cover and light to no wind. They begin to form as the sun set and often continue into the morning. Their presence can be indicated by ground fog; however, if fog is not present, inversions can also be identified by the movement of smoke from a ground source or an aircraft smoke generator. Smoke that layers and moves laterally in a connected cloud (under low wind conditions) indicates an inversion, while smoke that moves upwards and rapidly dissipates indicates good vertical air mixing.

Sensitive Areas

The pesticide should only be applied when the potential for drift to adjacent sensitive areas (e.g., residential areas, bodies of water, known habitat for threatened or endangered species, non-target crops) is minimal (e.g. when wind is blowing away from the sensitive areas).

FOR AERIAL APPLICATION IN CALIFORNIA ONLY

EXTREME CARE MUST BE EXERCISED TO AVOID CONTACT OF SPRAY WITH FOLIAGE, GREEN STEMS, OR FRUIT OF DESIRABLE CROPS, PLANTS, TREES, OR OTHER DESIRABLE VEGETATION SINCE SEVERE DAMAGE OR DESTRUCTION MAY RESULT.

Noncrop Sites

When applied as directed and under the conditions described in the "Weeds Controlled" section of the label booklet for this product, this herbicide will control or partially control the labeled weeds growing in the following industrial, recreational and public areas, such as airports, ditch banks, dry ditches, dry canals, fencerows, golf courses, highways, industrial plant sites, lumber yards, manufacturing sites, office complexes, parking areas, parks, petroleum tank farms and pumping installations, pipelines, power and telephone rights-of-way, railroads, roadsides (guiderails, shoulders), schools, storage areas, utility substations, warehouse areas and other public areas.

AVOID DRIFT - DO NOT APPLY WHEN WINDS ARE GUSTY OR UNDER ANY OTHER CONDITION WHICH WILL ALLOW DRIFT. DRIFT MAY CAUSE DAMAGE TO ANY VEGETATION CONTACTED TO WHICH TREATMENT IS NOT INTENDED. TO PREVENT INJURY TO ADJACENT DESIRABLE VEGETATION, APPROPRIATE BUFFER ZONES MUST BE MAINTAINED.

Use the following guidelines when aerial applications are made near crops or desirable perennial vegetation after bud break and before total leaf drop, and/or near other desirable vegetation or annual crops.

1. Do not apply within 100 feet of all desirable vegetation or crop(s).
2. If wind up to 5 miles per hour is blowing toward desirable vegetation or crop(s), do not apply within 500 feet of the desirable vegetation or crop(s).
3. Winds blowing from 5 to 10 miles per hour toward desirable vegetation or crop(s) may require buffer zones in excess of 500 feet.
4. Do not apply when winds are in excess of 10 miles per hour or when inversion conditions exist.

Coarse sprays are less likely to drift; therefore, do not use nozzles or nozzle configurations which dispense spray as fine spray droplets.

Do not angle nozzles forward into the airstream and do not increase spray volume by increasing nozzle pressure above the manufacturer's recommendation.

Drift control additives may be used. When a drift control additive is used, read and carefully observe the cautionary statements and all other information appearing on the additive label.

Ensure uniform application – to avoid streaked, uneven or overlapped application, use appropriate marking devices.

Thoroughly wash aircraft, especially landing gear, after each day of spraying to remove residues of this product accumulated during spraying or from spills. **PROLONGED EXPOSURE OF THIS PRODUCT TO UNCOATED STEEL SURFACES MAY RESULT IN CORROSION AND POSSIBLE FAILURE OF THE PART. LANDING GEAR IS MOST SUSCEPTIBLE.** The maintenance of an organic coating (paint) which meets aerospace specification MIL-C-38413 may prevent corrosion.

FOR AERIAL APPLICATION IN FRESNO COUNTY, CALIFORNIA ONLY

From February 15 through March 31 only. For aerial application outside of these dates, refer to the "FOR AERIAL APPLICATION IN CALIFORNIA ONLY" section printed above.

APPLICABLE AREA

This supplement only applies to the area contained inside the following boundaries within Fresno County, California only.

North: Fresno County line

South: Fresno County line

East: State Highway 99

West: Fresno County line

GENERAL INFORMATION

Always read and follow the label directions and precautionary statements for all products used in the aerial application.

Observe the following directions to minimize off-site movement during aerial application of this herbicide. Minimization of off-site movement is the responsibility of the grower, Pest Control Advisor, and aerial applicator.

WRITTEN RECOMMENDATIONS

A written recommendation **MUST** be submitted by or on behalf of the applicator to the Fresno County Agricultural Commissioner 24 hours prior to the application. This written recommendation **MUST** state the proximity of surrounding crops, and that conditions of each manufacturer's applicable product label(s) and this label have been satisfied.

AERIAL APPLICATOR TRAINING AND EQUIPMENT

Aerial application of this herbicide is limited to pilots who have successfully completed a Fresno County Agricultural Commissioner and California Department of Pesticide Regulation approved training program for aerial application of herbicides. All aircraft must be inspected, critiqued in flight, and certified at a Fresno County Agricultural Commissioner approved fly-in. Test and calibrate spray equipment at intervals sufficient to insure that proper rates of herbicides and adjuvants are being applied during commercial use. Applicator must document such calibrations and testing. Demonstration of performance at Fresno County Agricultural Commissioner approved "fly-ins" constitutes such documentation, or other written records showing calculations and measurements of flight and spray parameters acceptable to the Fresno County Agricultural Commissioner.

Application at night – Do not apply this product by air earlier than 30 minutes prior to sunrise and/or later than 30 minutes after sunset without prior permission from the Fresno County Agricultural Commissioner.

To report known or suspected misuse of this product or for additional information on the proper aerial application of this product, call 1-800-852-5234.

Read the "WARRANTY" section in this label booklet before using this product.

BROADCAST EQUIPMENT

For control of annual or perennial weeds listed on this label using broadcast equipment - Use the recommended rates of this product per acre as a broadcast spray unless otherwise specified on this label. See the "WEEDS CONTROLLED" section of this label for specific rates.

CONTROLLED DROPLET APPLICATION (CDA)

Apply these spray solutions in properly maintained and calibrated equipment capable of delivering desired volumes.

The rate of this product applied per acre by vehicle-mounted CDA equipment must not be less than the amount recommended in this label when applied by conventional broadcast equipment.

For vehicle-mounted CDA equipment apply [3 to 15 gallons of water per acre] or [in a sufficient volume of water to assure thorough coverage.]

For the control of labeled annual weeds with hand-held CDA units, apply a 20 percent solution. For the control of labeled perennial weeds, apply a 20 to 40 percent solution of this product.

Controlled droplet application equipment produces a spray pattern which is not easily visible. Extreme care must be exercised to avoid spray or drift contacting the foliage or any other green tissue of desirable vegetation, as damage or destruction may result.

HAND-HELD AND HIGH-VOLUME EQUIPMENT

Use Course Sprays Only

Mix this product in clean water and apply to foliage of vegetation to be controlled. For applications made on a spray-to-wet basis, spray coverage should be uniform and complete. Do spray to the point of runoff.

For control of annual weeds listed on this label, apply a 0.5 percent solution of this product plus nonionic surfactant to weeds less than 6 inches in height or runner length. Apply prior to seedhead formation in grass or bud formation in broadleaf weeds. For agricultural uses, allow 3 or more days before tillage or mowing.

For annual weeds over 6 inches tall, or when not using additional surfactant, or unless otherwise specified, use a 1 percent solution. For best results, use a 2 percent solution on harder-to-control [perennials, such as Bermudagrass, Dock, Field bindweed, Hemp dogbane, Milkweed and Canada thistle.] or [weeds.]

When using application methods which result in less than complete coverage, use a 5 percent solution for annual and perennial weeds and a 5 to 10 percent solution for woody brush.

See the "MIXING FOR HAND-HELD SPRAYERS" section of this label for specific rates.

SELECTIVE EQUIPMENT

This product may be applied through a recirculating spray system, shielded and hooded applicators, or wiper applicators after dilution and thorough mixing with water to listed weeds growing in any noncrop site specified on this label and only when specifically recommended in cropping systems.

AVOID CONTACT WITH DESIRABLE VEGETATION

Contact of the herbicide solution with the desirable vegetation may result in damage or destruction. Applicators used above desired vegetation should be adjusted so that the lowest spray stream or wiper contact point is at least 2 inches above the desirable vegetation. Droplets, mist, foam, or splatter of the herbicide solution settling on desirable vegetation may result in discoloration, stunting or destruction.

Applications made above the crops should be made when the weeds are a minimum of 6 inches above the desirable vegetation. Better results may be obtained when more of the weed is exposed to the herbicide solution. Weeds not contacted by the herbicide solution will not be affected. This may occur in dense clumps, severe infestations or when the height of the weeds varies so that not all weeds are contacted. In these instances, repeat treatment may be necessary.

SHIELDED AND HOODED APPLICATORS

When applied as directed under conditions described for shielded applicators, this product will control those weeds listed in the "WEEDS CONTROLLED" section of this label. Use the following equation to convert from a broadcast rate per acre to a band rate

$$\frac{\text{Band width in inches}}{\text{Row width in inches}} \times \text{Herbicide Broadcast RATE per acre} = \text{Herbicide Band RATE per acre}$$

$$\frac{\text{Band width in inches}}{\text{Row width in inches}} \times \text{Broadcast VOLUME of solution per acre} = \text{Band VOLUME of solution per acre}$$

Use nozzles that provide uniform coverage within the treated area. Keep shields on shielded sprayers adjusted to protect desirable vegetation. **EXTREME CARE MUST BE EXERCISED TO AVOID CONTACT WITH DESIRABLE VEGETATION.**

A hooded sprayer is a type of shielded applicator. The spray pattern is completely enclosed on the top and all 4 sides by a hood, thereby shielding the crop from the spray solution. This equipment must be set up and operated in a manner that avoids bouncing or raising the hoods off the ground in any way. If the hoods are raised, spray particles may escape and come into contact with the crop, causing damage or destruction of the crop. The spray hoods must be operated on the ground or skimming across the ground. Tractor speed must be adjusted to avoid bouncing of the spray hoods. Avoid operation on rough or sloping ground where the spray hoods might be raised off the ground.

For specific rates of application and instructions for control of various annual and perennial weeds, see the "WEEDS CONTROLLED" section of this label.

WIPER APPLICATORS

Wiper applicators are devices that physically wipe appropriate amounts of this product directly onto the weed.

Equipment must be designed, maintained and operated to prevent the herbicide solution from contacting desirable vegetation. Operate this equipment at ground speeds no greater than 5 mph. Performance may be improved by reducing speed in areas of heavy weed infestations to ensure adequate wiper saturation. Better results may be obtained if 2 applications are made in opposite directions.

Avoid leakage or dripping onto desirable vegetation. Adjust height of applicator to ensure adequate contact with weeds. Keep wiping surfaces clean. Be aware that, on sloping ground, the herbicide solution may migrate, causing dripping on the lower end and drying of the wicks on the upper end of a wiper applicator.

Do not use wiper equipment when weeds are wet.

Mix only the amount of solution to be used during a 1-day period, as reduced activity may result from use of leftover solutions. Clean wiper parts immediately after using this product by thoroughly flushing with water.

Do not add surfactant to the herbicide solution for wiper applications.

For Rope or Sponge Wick-Applicators - Mix 1 gallon of this product in 2 gallons of water to prepare a 33 percent solution. Apply this solution to weeds listed in this "WIPER APPLICATORS" section.

For Porous-Plastic Applicators - Solutions ranging from 33 to 100 percent of this product in water may be used in porous-plastic wiper applicators.

When applied as recommended under the conditions described for "WIPER APPLICATORS", this product CONTROLS the following weeds.

ANNUAL GRASSES

Corn	Panicum, Texas	Rye, common	Shattercane
------	----------------	-------------	-------------

ANNUAL BROADLEAVES

Sickelpod	Spanishneedles	Starbur, bristly
-----------	----------------	------------------

When applied as recommended under the conditions described for "WIPER APPLICATORS", this product SUPPRESSES the following weeds:

ANNUAL BROADLEAVES

Beggarweed, Florida	Pigweed, redroot	Ragweed, giant	Thistle, musk
Dogfennel	Ragweed, common	Sunflower	Velvetleaf

PERENNIAL GRASSES

Bermudagrass	Johnsongrass	Smutgrass	Vaseygrass
Guineagrass			

PERENNIAL BROADLEAVES

Dogbane	Milkweed	Nightshade, silverleaf	Thistle, Canada
Hemp			

WEEDS CONTROLLED

This herbicide controls many annual and perennial grasses and broadleaf weeds.

APPLICATIONS RATE TABLE

For annual and perennial weeds and woody brush

METHOD OF APPLICATION	APPLICATION RATE	RECOMMENDED SPRAY VOLUME (Gallons/Acre)
Broadcast		
Aerial	4 fluid ounces to	5 to 30
Ground	10 quarts per acre	3 to 100
Spray-to-Wet	1 to 2 percent	spray-to-wet
Handgun, Backpack, Mistblower	by volume	
Low Volume Directed Spray*		partial coverage
Handgun	5 to 10 percent	
Backpack	by volume	

*For low volume directed spray applications, coverage should be uniform with at least 50 percent of the foliage contacted. For best results, coverage of the top one-half of the plant is important.

NON-AGRICULTURAL USES

When applied as recommended under the conditions described, this product will control the following annual weeds. See "APPLICATION RATES TABLE" for recommended spray volumes.

Barley	Crabgrass	Horseweed/Marestail	Pigweed, (Redroot, Smooth)	Shepherdspurse
Barnyardgrass	Dwarfandelion	Johnsongrass,	Primrose, cutleaf evening	Sicklepod
Bluegrass,	Falseflax, smallseed	seedling	Pusley, Florida	Signalgrass, broadleaf
(Annual, Bulbous)	Filaree	Lambsquarters,	Rice, red	Spanishneedles
Brome, downy	Foxtail	common	Rocket, London	Sprangletop
Buttercup	Foxtail, Carolina	Morningglory	Rye	Spurge, annual
Cheatgrass	Geranium, Carolina	Mustard, (Blue, Tansy,	Ryegrass, Italian	Spurry, umbrella
Chickenweed	Goatgrass, jointed	Tumble, Wild)	Sandbur, field	Stinkgrass
(Common, Mouseear)	Goosegrass	Oats, wild	Shattercane	Teaweed
Cocklebur	Groundsel, common	Panicum, (tall, Texas)		Wheat
Corn	Henbit	Pennycress, field		Witchgrass

AGRICULTURAL USES

Apply to actively growing grass and broadleaf weeds. To prevent seed production, applications should be made prior to seedhead formation. This product does not provide residual control; therefore, delay application until maximum weed emergence. Repeat treatments may be necessary to control later germinating weeds. For maximum agronomic benefit, apply when weeds are 6 inches or less in height. Allow at least 3 days after treatment before tillage.

LOW-VOLUME BROADCAST APPLICATION (LOW-RATE TECHNOLOGY)

When applied as directed under the conditions described, this product will control the annual weeds listed below in the "ANNUAL WEEDS RATE TABLE" when:

1. Apply in sufficient volume of water to ensure thorough coverage.

NOTE

- Do not tank-mix with soil residual herbicides when using these rates unless otherwise specified.
- For weeds that have been mowed, grazed or cut, allow regrowth to occur prior to treatment.
- Refer to the "TANK MIXTURES" portion of this section for control of additional broadleaf weeds.

ANNUAL WEEDS RATE TABLE

WEED SPECIES	MAXIMUM HEIGHT/LENGTH (Inches)	USE RATE (Fluid Ounces/Acre)
Barley	12	12
Barnyardgrass	6 [0 to 4 4 to 6]	12 16 ¹ 24 ¹
Bluegrass, bulbous	6	16
Bluegrass, annual	6	12
Brome, downy**	6	12
Buttercup	12	16
Cheat	6	16
Chickweed, common	6	16
Chickweed, mouseear	6	16
Cocklebur	12	16
Corn	6	16
Crabgrass	12	16
Dwarf dandelion	12	16
Falseflax, smallseed	12	16
Filaree	12	48
Foxtail	12	8
Foxtail, Carolina	12	16
Geranium, Carolina	12	32
Goatgrass, jointed	6	16
Goosegrass	12	32
Groundsel, common	6	16
Henbit	6	16
Horseweed/Marestail	6 7 to 12	16 24
Johnsongrass, seedling	12	16

(continued)

ANNUAL WEEDS RATE TABLE (continued)

WEED SPECIES	MAXIMUM HEIGHT/LENGTH (Inches)	USE RATE (Fluid Ounces/Acre)
Lambsquarters, common	6 7 to 12	12 24
Morningglory	2	16
Mustard, blue	6	12
Mustard, tansy	6	12
Mustard, tumble	6	12
Mustard, wild	6	12
Oats, wild	12	16
Panicum, fall	12	16
Panicum, Texas	12	16
Pennycress, field	6	16
Pigweed, redroot	12	16
Pigweed, smooth	12	16
Primrose, cutleaf evening	12	32
Pusley, Florida	12	32
Rice, red	4	32
Rocket, London	6	16
Rye	12	12
Ryegrass, Italian	6	16
Shattercane	12	12
Shepherdspurse	6	16
Sicklepod	2 3 to 4 5 to 12	16 24 32
Signalgrass, broadleaf	4	24
Spanishneedles	5 to 12	32
Sprangletop	6 12	32 48
Spurge, annual	6 7 to 12	16 24
Spurry, umbrella	6	12
Stinkgrass	12	12
Teaweed	4	32
Wheat	18	12
Witchgrass	12	16

¹ Use these rates to control barnyardgrass in Alabama, Arkansas, Mississippi, Missouri, Louisiana and Texas for preplant treatments.

*For those rates less than 32 fluid ounces per are, this product at rates up to 32 fluid ounces per acre may be used where heavy weed densities exist.

**For control in no-till systems, use 16 fluid ounces per acre.

HIGH-VOLUME BROADCAST APPLICATIONS

When applied as directed under the conditions described, this product will control the weeds listed below when water carrier volumes are 10 gallons or more for sufficient coverage. Apply 1 to 1.5 quarts of this product per acre plus additional surfactant according to the manufacturers rates and recommendations for of the surfactant. Use 1 quart per acre if weeds are less than 6 inches tall and 1.5 quarts per acre if weeds are over 6 inches tall. If weeds have been mowed, grazed or cut, allow adequate time for new growth to reach recommended stages prior to treatment. These rates will also provide control of weeds listed in the "LOW-VOLUME BROADCAST APPLICATION" section.

Balsamapple*	Fleabane, hairy	Panicum	Sowthistle, annual
Bassia, fivehook	Fleabane	Ragweed, common	Sunflower
Brome	Kochia	Ragweed, giant	Thistle, Russian
Fiddleneck	Lettuce, prickly	Smartweed, Pennsylvania	Velvetleaf.

*Apply with hand-held equipment only.

PERENNIAL WEEDS

Apply this product as follows to control or destroy most perennial weeds:

When applied as recommended under the conditions described, this product WILL CONTROL the following PERENNIAL WEEDS:

Alfalfa	Cogongrass	Nightshade, silverleaf	Starthistle, yellow
Alligatorweed*	Dallisgrass	Nutsedge (Purple, Yellow)	Sweet potato, wild*
Anise (fennel)	Dandelion	Orchardgrass	Thistle, Canada
Artichoke, Jerusalem	Dock, curly	Pampasgrass	Thistle, artichoke
Bahiagrass	Dogbane, hemp	Paragrass	Timothy
Bentgrass	Fescues	Phragmites*	Torpedograss*,
Bermudagrass	Fescue, tall	Poison hemlock	Trumpetcreep*
Bermudagrass, water	Guineagrass	Quackgrass	Vaseygrass
(knotgrass)	Horsenettle	Redvine*	Velvetgrass
Bindweed, field	Horseradish	Reed, giant	Wheatgrass, western
Bluegrass, Kentucky	Ice plant	Ryegrass, perennial	
Cattail	Mullein, common	Smartweed, swamp	
Clover (Red, White),	Napiergrass	Spurge, leafy*	

*Partial control.

This product is not registered in California for use on water Bermudagrass.

PERENNIAL WEEDS RATE TABLE

Apply to actively growing perennial weeds. See "APPLICATION RATES TABLE" for recommended spray volumes. For other perennials listed on this label and not found in the table below, apply 3 to 5 quarts of this product per acre. Apply when actively growing and most have reached early head or early bud stage of growth. For agricultural uses, allow 7 or more days after application before tillage.

Additional surfactant may be used. If additional surfactant is to be used follow the manufacturers rates and recommendations for use of the surfactant.

PERENNIAL WEEDS	USE RATE Broadcast Spray (Quarts/Acre)	USE RATE Hand Held Equipment (Percent Solution)
Alfalfa	1	
	Make application after the last hay cutting in the fall. Allow alfalfa to regrow to a height of 6 to 8 inches or more prior to treatment. For agricultural uses applications should be followed with deep tillage at least 7 days after treatment, but before soil freeze-up.	
Alligatorweed	4	1.5
	Partial control. Apply when most of the plants are in bloom. Repeat applications will be required to maintain such control.	
Anise (fennel)	--	1 to 2
	Apply as a spray-to-wet treatment. Optimum results are obtained when plants are treated at the bud to full-bloom stage of growth. Repeat applications may be needed in succeeding years to control plants arising from seeds.	
Artichoke, Jerusalem	3 to 5	
	Apply when actively growing and most have reached early head or early bud stage of growth. For agricultural uses allow 7 or more days after application before tillage.	

PERENNIAL WEEDS RATE TABLE (continued)

PERENNIAL WEEDS	USE RATE Broadcast Spray (Quarts/Acre)	USE RATE Hand Held Equipment (Percent Solution)
Bahiagrass	3 to 5	
	Apply when actively growing and most have reached early head or early bud stage of growth. For agricultural uses allow 7 or more days after application before tillage.	
Bentgrass	1.5	
	For suppression in grass seed production areas. For ground applications only. Ensure entire crown area has resumed growth prior to a fall application. Bentgrass should be actively growing and have at least 3 inches of growth. Tillage prior to treatment should be avoided. Tillage 7 to 10 days after application is recommended for best results. Failure to use tillage after treatment may result in unacceptable control.	
Bermudagrass	3 to 5	--
	For control, apply 5 quarts of this product per acre. For partial control, apply 3 quarts per acre. Treat when Bermudagrass is actively growing and seedheads are present. Retreatment may be necessary to maintain control. Allow 7 or more days after application before tillage.	
Bermudagrass, water (knotgrass)	1 to 1.5	--
	<p>Apply when water Bermudagrass is actively growing and 12 to 18 inches in length.</p> <p>For agricultural uses apply 1.5 quarts of this product per acre. Allow 7 or more days before tilling, flushing or flooding the field.</p> <p>Fall applications only: - Fallow field should be tilled prior to application. Apply prior to frost on water Bermudagrass that is actively growing and 12 to 18 inches in length.</p> <p>For agricultural uses apply 1 quart of this product per acre Allow 7 or more days before tillage.</p>	
Bindweed, field	1 to 5	--
	<p>For control, apply 4 to 5 quarts of this product per acre west of the Mississippi River and 3 to 4 quarts east of the Mississippi River. Apply when the weeds are actively growing and are at or beyond full bloom. Do not treat when weed is under drought stress as good soil moisture is necessary for active growth. For best results, apply in late summer or fall. Fall treatments must be applied before a killing frost. Allow 7 or more days after application before tillage.</p> <p>Also for control, apply 2 quarts of this product plus 0.5 pound a.i. of Dicamba per acre. At these rates, apply using ground application only.</p> <p>Agricultural Uses: The following tank mixtures with 2,4-D may be applied using aerial application equipment (except in California) in fallow and reduced tillage systems only.</p> <p>For suppression on irrigated agricultural land, apply 1 to 2 quarts of this product plus 1 pound a.i. of 2,4-D per acre with ground equipment only. Applications should be made following harvest or in fall fallow ground when the bindweed is actively growing and the majority of runners are 12 inches or more in length. The use of at least one irrigation will promote active bindweed growth.</p> <p>For suppression, apply 16 fluid ounces of this product plus 0.5 pound a.i. of 2,4-D per acre in 3 to 10 gallons of water per acre for ground applications and 3 to 5 gallons of water per acre for aerial applications.</p> <p>Applications should be delayed until maximum emergence has occurred and when vines are between 6 to 18 inches in length.</p> <p>In California only, apply 1 to 5 quarts of this product per acre. Actual rate needed for suppression or control will vary within this range depending on local conditions. For suppression on irrigated land where annual tillage is performed, apply 1 quart of this product per acre plus surfactant. Apply to actively growing bindweed that has reached a length of 12 inches or greater. Allow maximum weed emergence and runner growth. Do not treat when weeds are under drought stress as good soil moisture is necessary for active growth. For agricultural uses allow 3 or more days after application before tillage.</p>	

PERENNIAL WEEDS RATE TABLE (continued)

PERENNIAL WEEDS	USE RATE Broadcast Spray (Quarts/Acre)	USE RATE Hand Held Equipment (Percent Solution)
Bluegrass, Kentucky	2	--
	Apply when the grasses are actively growing and most plants have reached boot-to-early seedhead stage of development. For partial control in pasture or hay crop renovation, apply 1 to 1.5 quarts of this product per acre. Apply to actively growing plants when most have reached 4 to 12 inches in height. Allow 7 or more days after application before tillage.	
Blueweed, Texas	3 to 5	--
	Apply 4 to 5 quarts of this product per acre west of the Mississippi River and 3 to 4 quarts per acre east of the Mississippi River. Apply when weed is actively growing and is at or beyond full bloom. Do not treat when weed is under drought stress as good soil moisture is necessary for active growth. New leaf development indicates active growth. For best results, apply in late summer or fall. Fall treatments must be applied before a killing frost. Allow 7 or more days after application before tillage.	
Brackenfern	3 to 4	1 to 1.5
	Apply to fully expanded fronds which are at least 18 inches long.	
Bromegrass, smooth	2	--
	Apply when the grasses are actively growing and most plants have reached boot-to-early seedhead stage of development. For partial control in pasture or hay crop renovation, apply 1 to 1.5 quarts of this product per acre. For agricultural uses apply to actively growing plants when most have reached 4 to 12 inches in height. and Allow 7 or more days after application before tillage.	
Bursage, woollyleaf	1 to 2 plus 1 pint Dicamba	--
	For control, apply 2 quarts of this product plus 1 pint of Dicamba per acre. For partial control, apply 1 quart of this product plus 1 pint of Dicamba per acre. Apply when plants are producing new active growth which has been initiated by moisture for a least 2 weeks and when plants are at or beyond flowering.	
Canarygrass, reed	2 to 3	--
	For best results, apply to actively growing plants when most have reached the boot-to-head stage of growth. Allow 7 or more days after application before tillage.	
Cattail	3 to 5	--
	Apply when actively growing and most have reached early head to early bud stage of growth. Allow 7 or more days after application before tillage.	
Clover: Red White	3 to 5	--
	Apply when actively growing and most have reach early head or early bud stage of growth. Allow 7 or more days after application before tillage.	
Cogongrass	3 to 5	
	Apply when Cogongrass is at least 18 inches tall and actively growing in late summer or fall. Due to uneven stages of growth and the dense nature of vegetation preventing good spray coverage, repeat treatments may be necessary to maintain control. Allow 7 or more days after application before tillage or mowing.	

(continued)

PERENNIAL WEEDS RATE TABLE (continued)

PERENNIAL WEEDS	USE RATE Broadcast Spray (Quarts/Acre)	USE RATE Hand Held Equipment (Percent Solution)
Dallisgrass	3 to 5	--
	Apply when actively growing and most have reached early head or early bud stage of growth. Allow 7 or more days after application before tillage.	
Dandelion	3 to 5	--
	Apply when plants are actively growing and most have reached the early bud stage of growth. Allow 7 or more days after application before tillage. Also for control, apply 16 fluid ounces of this product plus 0.5 pound a.i. 2,4-D per acre.	
Dock, curly	3 to 5	--
	Apply when plants are actively growing and most have reached the early bud stage of growth. Allow 7 or more days after application before tillage. Also for control, apply 16 fluid ounces of this product plus 0.5 pound a.i. 2,4-D per acre.	
Dogbane, hemp	4	--
	Apply when actively growing and when most weeds have reached the late bud to flower stage of growth. Following crop harvest or mowing, allow weeds to regrow to a mature stage prior to treatment. For best results, apply in late summer or fall. Allow 7 or more days after application before tillage. For suppression, apply 16 fluid ounces of this product plus 0.5 pound a.i. of 2,4-D per acre. Delay applications until maximum emergence of dogbane has occurred.	
Fescue (except tall)	3 to 5	--
	Apply when actively growing and most have reached early head or early bud stage of growth. Allow 7 or more days after application before tillage.	
Fescue, tall	1 to 5	--
	Apply 3 quarts of this product in per acre to actively growing plants when most have reached boot-to-early seedhead stage of development. Fall applications only: Apply 1 quart of this product per acre. Apply to fescue in the fall when actively growing and plants have 6 to 12 inches of new growth. A sequential application of 1 pint per acre of this product plus nonionic surfactant will improve long-term control and control seedlings germinating after fall treatments or the following spring. Allow 7 or more days after application before tillage.	
Guineagrass	3	1
	Apply to actively growing guineagrass when most has reached at least the 7-leaf stage of growth. Ensure thorough coverage when using hand-held equipment. Allow 7 or more days after application before tillage.	
Horsenettle	3 to 5	--
	Apply when actively growing and most have reached early head or early bud stage of growth. Allow 7 or more days after application before tillage.	
Horseradish	4	--
	Apply when actively growing and when most weeds have reached the late bud to flower stage of growth. Following crop harvest or mowing, allow weeds to regrow to a mature stage prior to treatment. For best results, apply in late summer or fall. Allow 7 or more days after application before tillage.	

(continued)

PERENNIAL WEEDS RATE TABLE (continued)

PERENNIAL WEEDS	USE RATE Broadcast Spray (Quarts/Acre)	USE RATE Hand Held Equipment (Percent Solution)
Ice plant	--	1.5 to 2
	Apply to plants that are actively growing. Thorough coverage is necessary for best control.	
Johnsongrass	0.5 to 3	1
	<p>In noncrop, or areas where annual tillage (no-till) is not performed, apply 2 to 3 quarts of this product acre. In annual cropping systems apply 1 to 2 quarts of this product per acre. Use 2 quarts of this product when applying 10 to 40 gallons of water per acre. For best results, apply to actively growing plants when most have reached the boot-to-head stage of growth or in the fall prior to frost. Do not tank-mix with residual herbicides when using the 1 quart per acre rate. Allow 7 or more days after application before tillage.</p> <p>For burndown of Johnsongrass, apply 1 pint per acre before the plants reach a height of 12 inches. For this use, allow at least 3 days after treatment before tillage.</p> <p>Spot treatment (partial control or suppression)-Apply a 1 percent solution of this product when Johnsongrass is 12 to 18 inches in height. Coverage should be uniform and complete.</p>	
Kikuyugrass	2 to 3	--
	Spray when most kikuyugrass is at least 8 inches in height (3 or 4-leaf stage of growth) and actively growing. Allow 3 or more days after application before tillage.	
Knapweed	2 to 3	--
	Apply when actively growing and when most weeds have reached the late bud to flower stage of growth. Following crop harvest or mowing, allow weeds to regrow to a mature stage prior to treatment. For best results, apply in late summer or fall. Allow 7 or more days after application before tillage.	
Lantana	--	1 to 1.25
	Apply to actively growing lantana at or beyond the bloom stage of growth. Use the higher application rate for plants that have reached the woody stage of growth. Allow 7 or more days after application before tillage.	
Lespedeza	3 to 5	--
	Apply when actively growing and most have reached early head or early bud stage of growth. Allow 7 or more days after application before tillage.	
Milkweed, common	3	--
	Apply when actively growing and most of the milkweed has reached the late bud to flower stage of growth. Following small grain harvest or mowing, allow milkweed to regrow to a mature stage prior to treatment. Allow 7 or more days after application before tillage.	
Muhly, wirestem	1 to 2	--
	Use 1 quart of this product per acre. Use 2 quarts per acre in pasture, sod, or noncrop areas. Spray when the wirestem muhly is 8 inches or more in height and actively growing. Do not till between harvest and fall applications or in the fall or spring prior to spring applications. This product will not provide residual control of wirestem muhly from seeds which germinate after application of this product. Do not tank mix with residual herbicides when using the 1 quart per acre rate. Allow 3 or more days after application before tillage.	

(continued)

PERENNIAL WEEDS RATE TABLE (continued)

PERENNIAL WEEDS	USE RATE Broadcast Spray (Quarts/Acre)	USE RATE Hand Held Equipment (Percent Solution)
Mullein, common	3 to 5	--
	Apply when actively growing and most have reached early head or early bud stage of growth. Allow 7 or more days after application before tillage.	
Napiergrass	3 to 5	--
	Apply when actively growing and most have reached early head or early bud stage of growth. Allow 7 or more days after application before tillage.	
Nightshade, silverleaf	2	--
	For control, applications should be made when at least 60 percent of the plants have berries. Fall treatments must be applied before a killing frost. Do not treat when weed is under drought stress as good soil moisture is necessary for active growth. Allow 7 or more days after application before tillage.	
Nutsedge: Purple Yellow	0.5 to 3	1 to 2
	<p>Apply 3 quarts of this product per acre as a broadcast spray, or apply a 1 to 2 percent solution from hand-held equipment to control existing nutsedge plants and immature nutlets attached to treated plants. Treat when plants are in flower or when new nutlets can be found at rhizome tips. Nutlets which have not germinated will not be controlled and may germinate following treatment. Repeat treatments will be required for longterm control of ungerminated tubers.</p> <p>Sequential applications of 1 to 2 quarts of this product per acre will provide control. Make applications when a majority of the plants are in the 3 to 5-leaf stage (less than 6 inches tall). Repeat this application, as necessary, when newly emerging plants reach the 3 to 5-leaf stage. Subsequent applications will be necessary for long-term control.</p> <p>For suppression to partial control of existing plants, apply 1 pint to 2 quarts of this product per acre. Treat when plants have 3 to 5 leaves and most are less than 6 inches tall. Repeat treatments will be required to control subsequent emerging plants or regrowth of existing plants. Wait 7 days after treatment before tillage or mowing.</p>	
Orchardgrass	1 to 2	--
	<p>Apply 2 quarts of this product when the grasses are actively growing and most plants have reached boot-to-early seedhead stage of development. For partial control in pasture or hay crop renovation, apply 1 to 1.5 quarts of this product per acre. Apply to actively growing plants when most have reached 4 to 12 inches in height. Allow 7 or more days after application before tillage.</p> <p>Sods going to no-till corn: Apply 1 to 1.5 quarts of this product per acre. Apply to orchardgrass that is a minimum of 12 inches tall for spring applications and 6 inches tall for fall applications. Allow at least 3 days following application before planting. A sequential application of atrazine will be necessary for optimum results.</p>	
Pampasgrass	--	1.5 to 2
	Apply to plants that are actively growing. Pampasgrass should be at or beyond the boot stage of growth. Thorough coverage is necessary for best control.	
Paragrass	3 to 5	--
	Apply when actively growing and most have reached early head or early bud stage of growth. Allow 7 or more days after application before tillage.	

(continued)

PERENNIAL WEEDS RATE TABLE (continued)

PERENNIAL WEEDS	USE RATE Broadcast Spray (Quarts/Acre)	USE RATE Hand Held Equipment (Percent Solution)
Phragmites	3 to 5	1 to 2
	For partial control of phragmites in Florida and the counties of other states bordering the Gulf of Mexico, apply 5 quarts per acre as a broadcast spray or apply a 2 percent solution from hand-held equipment. In other areas of the U.S., apply 3 quarts per acre as a broadcast spray or apply a 1 percent solution from hand-held equipment for partial control. For best results, treat during late summer or fall months or when plants are actively growing and in full bloom. Treatment before or after this stage may lead to reduced control. Due to the dense nature of the vegetation, which may prevent good spray coverage or uneven stages of growth, repeat treatments may be necessary to maintain control. Visual control symptoms will be slow to develop.	
Poison hemlock	--	1 to 2
	Apply as a spray-to-wet treatment. Optimum results are obtained when plants are treated at the bud to full-bloom stage of growth. Repeat applications may be needed in succeeding years to control plants arising from seeds.	
Quackgrass	1 to 2	--
	<p>In Annual Cropping Systems, or in Pastures and Sods Followed by Deep Tillage: Apply 1 to 2 quarts of this product per acre. Do not tank mix with residual herbicides when using the 1 quart rate. Spray when quackgrass is 6 to 8 inches in height and actively growing. Do not till between harvest and fall applications or in fall or spring prior to spring application. In pastures or sods, for best results use a moldboard plow. Allow 3 or more days after application before tillage.</p> <p>Pasture or Sod or Other Noncrop Areas Where Deep Tillage is Not Planned Following Application: Apply 2 to 3 quarts per acre. Spray when the quackgrass is greater than 8 inches tall and actively growing. Do not till between harvest and fall application or in fall or spring prior to spring application. Allow 3 or more days after application before tillage.</p>	
Redvine	3/4 to 2	--
	For suppression, apply 24 fluid ounces of this product per acre at each of two applications 7 to 14 days apart or a single application of 2 quarts per acre. Apply in late September or early October to actively growing plants, which are at least 18 inches tall and have been growing 45 to 60 days since the last tillage operation. Make applications at least 1 week before a killing frost.	
Reed, giant	--	2
	For control. Apply when plants are actively growing. Best results are obtained when applications are made in late summer to fall.	
Ryegrass, perennial	1 to 3	1
	In noncrop, or areas where annual tillage (no-till) is not performed, apply 2 to 3 quarts of this product per acre. In annual cropping systems apply 1 to 2 quarts of this product per acre. For best results, apply to actively growing plants when most have reached the boot-to-head stage of growth or in the fall prior to frost. Do not tank-mix with residual herbicides when using the 1 quart per acre rate. Allow 7 or more days after application before tillage.	

(continued)

PERENNIAL WEEDS RATE TABLE (continued)

PERENNIAL WEEDS	USE RATE Broadcast Spray (Quarts/Acre)	USE RATE Hand Held Equipment (Percent Solution)
Smartweed, swamp	3 to 5	--
	Apply 3 to 5 quarts of this product per acre when plants are actively growing and most have reached the early bud stage of growth. Allow 7 or more days after application before tillage. Also for control, apply 16 fluid ounces of this product plus 0.5 pound active ingredient of 2,4-D per acre in the late summer or fall. Apply when plants are actively growing and most have reached the early bud stage of growth. Allow 7 or more days after application before tillage.	
Spurge, leafy	1/2	--
	For suppression, apply 1/2 quart of this product plus 0.5 pound active ingredient 2,4-D per acre in the late summer or fall. Apply when plants are actively growing. If mowing has occurred prior to treatment, apply when most of the plants are 12 inches tall. Allow 7 or more days after application before tillage.	
Starthistle, yellow	2	2
	Best results are obtained when applications are made during periods of active growth, including the rosette, bolting and early flower stages.	
Sweet potato, wild	--	3
	Apply to actively growing weeds that are at or beyond the bloom stage of growth. Repeat applications may be required. Allow the plant to reach the recommended stage of growth before retreatment. Allow 7 or more days before tillage.	
Thistle, artichoke	--	2
	Apply to actively growing weeds that are at or beyond the bloom stage of growth. Repeat applications may be required. Allow the plant to reach the recommended stage of growth before retreatment. Allow 7 or more days before tillage.	
Thistle, Canada	1 to 3	--
	Apply 2 to 3 quarts of this product per acre. Apply to actively growing thistles when most are at or beyond the bud stage of growth. After harvest, mowing or tillage in the late summer or fall, allow at least 4 weeks for initiation of active growth and rosette development prior to the application of this product. Fall treatments must be applied before a killing frost. Allow 3 or more days after application before tillage. For suppression of Canada thistle, apply 1 quart per acre of this product, or 1 pint of this product plus 0.5 pound a.i. 2,4-D per acre, in the late summer or fall after harvest, mowing or tillage. Allow rosette regrowth to a minimum of 6 inches in diameter before treating. Applications can be made as long as leaves are still green and plants are actively growing at the time of application. Allow 3 or more days after application before tillage.	
Timothy	2 to 3	--
	For best results, apply to actively growing plants when most have reached the boot-to-head stage of growth. Allow 7 or more days after application before tillage.	
Torpedograss	4 to 5	--
	Partial control. Apply to actively growing torpedograss when most plants are at or beyond the seedhead stage of growth. Repeat application will be required to maintain control. Fall treatments must be applied before frost. Allow 7 or more days after application before tillage.	

PERENNIAL WEEDS RATE TABLE (continued)

PERENNIAL WEEDS	USE RATE Broadcast Spray (Quarts/Acre)	USE RATE Hand Held Equipment (Percent Solution)
Trumpet creeper	2	--
	For control. Apply to actively growing plants in late September or October, which are at least 18 inches tall and have been growing 45 to 60 days since the last tillage operation. Make applications at least 1 week before a killing frost.	
Vaseygrass	3 to 5	--
	Apply when actively growing and most have reach early head or early bud stage of growth. Allow 7 or more days after application before tillage.	
Velvetgrass	3 to 5	--
	Apply when actively growing and most have reached early head or early bud stage of growth. Allow 7 or more days after application before tillage.	
Wheatgrass, western	2 to 3	--
	For best results, apply to actively growing plants when most have reached the boot-to-head stage of growth. Allow 7 or more days after application before tillage.	

TANK MIXTURES

This product can be tank mixed with additional products to provide residual control of many of the listed hard to control problem species. The addition of recommended rates of 2,4-D or Dicamba based products will provide improved control of many annual and perennial weeds. This product may be tank-mixed with other herbicides provided the specific product is registered for use on these sites. When tank mixing, read and observe applicable use directions, precautions and limitations on the respective product labels. Use according to the most restrictive label directions for each product in the mixture.

WOODY BRUSH

When applied as recommended under the conditions described, this product CONTROLS or PARTIALLY CONTROLS the following woody brush, plants and trees:

Alder	Creepers, Virginia*	Madrone	Sagebrush, California
Ash	Dewberry	Manzanita	Salmonberry
Aspen, quaking	Dogwood*	Maple (red, sugar, vine)	Salt cedar
Bearmat (Bearclover)	Elderberry	Monkey Flower*	Sassafras
Beech	Elm*	Oak (Black*, Northern Pin,	Sourwood
Birch	Eucalyptus	Post, Red, Southern Red,	Sumac (Poison*, Smooth*,
Blackberry	Gorse	White*)	Winged*)
Blackgum	Hasardia*	Persimmon*	Sweetgum
Bracken	Hawthorn	Pine	Swordfern*
Broom (French, Scotch)	Hazel	Poison Ivy	Tallowtree, Chinese
Buckwheat, California*	Hickory*	Poison Oak	Tan Oak
Cascara*	Holly, Florida/Brazilian	Poplar, yellow*	Thimbleberry
Catsclaw*	Peppertree*	Raspberry	Tobacco, tree*
Ceanothus*	Honeysuckle	Redbud, eastern	Trumpet creeper
Chamise	Hornbeam, American*	Rose, multiflora	Waxmyrtle, southern*
Cherry (Bitter, Black, Pin),	Kudzu	Russian-olive***	Willow
Coyote brush	Locust, black*	Sage (Black, White)	

* Partial control

** See below for control or partial control instructions.

*** This product is not registered in California for use on Russian-olive.

NOTE: If brush has been mowed or tilled or trees have been cut, do not treat until regrowth has reached the recommended stages of growth. Apply this product when plants are actively growing and, unless otherwise directed, after full leaf expansion. Use the higher rate for larger plants and/or dense areas of growth. On vines, use the higher rate for plants that have reached the woody stage of growth. Best results are obtained when application is made in late summer or fall after fruit formation.

In arid areas, best results are obtained when application is made in the spring to early summer when brush species are at high moisture content and are flowering.

Ensure thorough coverage when using hand-held equipment. Symptoms may not appear prior to frost or senescence with fall treatments. Allow 7 or more days after application before tillage, mowing or removal. Repeat treatments may be necessary to control plants regenerating from underground parts or seed. Some autumn colors on undesirable deciduous species are acceptable provided no major leaf drop has occurred. Reduced performance may result if fall treatments are made following a frost.

See "DIRECTIONS FOR USE" and "MIXING, ADDITIVES and APPLICATION INSTRUCTIONS" sections of this label for labeled uses and specific application instructions.

WOODY BRUSH RATE TABLE

See "APPLICATION RATES TABLE" for recommended spray volumes.

For Other Woody Brush listed on this label but not found in the table below: For partial control, apply 2 to 5 quarts of this product per acre as a broadcast spray or as a 1 to 2 percent solution with hand-held equipment. For difficult to control perennial weeds and woody brush and trees, where plants are growing under stressed conditions or where infestations are dense, Razor may be used at 5 to 10 quarts per acre for enhanced results. **The annual maximum use rate for this product is 10.6 quarts per acre per year**

Additional surfactant may be used. If additional surfactant is to be used follow the manufacturers rates and recommendations for use of the surfactant.

WOODY BRUSH	USE RATE Broadcast Spray (Quarts/Acre)	USE RATE Hand Held Equipment (Percent Solution)
Alder	For control. 3 to 4	1 to 1.5
Ash	For partial control. 2 to 5	1 to 2
Aspen, quaking	For control. 2 to 3	1 to 1.5
Bearmat (bearclover)	For partial control. 2 to 5	1 to 2
Beech	For partial control. 2 to 5	1 to 2
Birch	For control. 2	1
Blackberry	For control. 3 to 4	1 to 1.5
	Make application after plants have reached full leaf maturity. Best results are obtained when applications are made in late summer or fall. After berries have set or dropped in late fall, blackberry can be controlled by applying a 3/4 percent solution of this product with hand-held equipment. For control of blackberries after leaf drop and until killing frost or as long as stems are green, apply 3 to 4 quarts of this product in 10 to 40 gallons of water per acre.	
Blackgum	For partial control. 2 to 5	1 to 2
Bracken	For partial control. 2 to 5	1 to 2
Broom: French and Scotch	For control.	1.5 to 2
Buckwheat, California	For partial control. 2 to 5	1 to 2
	Thorough coverage of foliage is necessary for best results.	
Cascara	For partial control. 2 to 5	1 to 2
Catsclaw	For partial control.	1 to 1.5
Ceanothus	For partial control. 2 to 5	1 to 2
Chamise	For control.	1
	Thorough coverage of foliage is necessary for best results.	
Cherry: Bitter, Black and Pin	For control. 2 to 3	1 to 1.5
Coyote brush	For control.	1.5 to 2
	Apply when at least 50 percent of the new leaves are fully developed.	
Creeper, Virginia	For partial control. 2 to 5	1 to 2
Dewberry	For control. 3 to 4	1 to 1.5

(continued)

WOODY BRUSH RATE TABLE (continued)

WOODY BRUSH	USE RATE Broadcast Spray (Quarts/Acre)	USE RATE Hand Held Equipment (Percent Solution)
Dogwood	For partial control. 2 to 5	1 to 2
Elderberry	For control. 2	1
Elm	For partial control. 2 to 5	1 to 2
Eucalyptus	For control. 2	
	Apply when eucalyptus resprouts are 6 to 12 feet tall. Ensure complete coverage. Apply when plants are growing actively. Avoid application to drought-stressed plants.	
Gorse	For partial control. 2 to 5	1 to 2
Hasardia	For partial control. 1 to 2	
	Thorough coverage of foliage is necessary for best results.	
Hawthorn	For control. 2 to 3	1 to 1.5
Hazel	For control. 2	1
Hickory	For partial control. 2 to 5	1 to 2
Holly, Florida/Brazilian Peppertree	For partial control. 2 to 5	1 to 2
Honeysuckle	For control. 3 to 4	1 to 1.5
Hornbeam, American	For partial control. 2 to 5	1 to 2
Kudzu	For control. 4	2
	Repeat applications will be required to maintain control.	
Locust, black	For partial control. 2 to 5	1 to 2
Madrone, resprouts	For suppression or partial control. 2	
	Apply to resprouts less than 3 to 6 feet tall. Best results are obtained with spring/early summer treatments.	
Manzanita	For partial control. 2 to 5	1 to 2
Maple, red	For control. 2 to 4	1 to 1.5
	Apply as a 1 to 1.5 percent solution when at least 50 percent of the new leaves are fully developed. For partial control, apply 2 to 4 quarts of this product per acre.	
Maple, sugar	For control. 1 to 1.5	
	Apply when at least 50 percent of the new leaves are fully developed.	
Maple, vine	For partial control. 2 to 5	1 to 2
Monkey flower	For partial control. 1 to 2	
	Thorough coverage of foliage is necessary for best results.	
Oak: Black and White	For partial control. 2 to 5	1 to 2
Oak: Northern Pin and Red	For control. 1 to 1.5	
	Apply when at least 50 percent of the new leaves are fully developed.	
Oak, post	For control. 3 to 4	1 to 1.5
Oak, southern red	For control. 2 to 3	1 to 1.5
Persimmon	For partial control. 2 to 5	1 to 2
Pine	For partial control. 2 to 5	1 to 2

(continued)

WOODY BRUSH RATE TABLE (continued)

WOODY BRUSH	USE RATE Broadcast Spray (Quarts/Acre)	USE RATE Hand Held Equipment (Percent Solution)
Poison Ivy	For control. 4 to 5	2
	Repeat applications may be required to maintain control. Fall treatments must be applied before leaves lose green color.	
Poison oak	For control. 4 to 5	2
	Repeat applications may be required to maintain control. Fall treatments must be applied before leaves lose green color.	
Poplar, yellow	For partial control. 2 to 5	1 to 2
Raspberry	For control. 3 to 4	1 to 1.5
Redbud, eastern	For partial control. 2 to 5	1 to 2
Rose, multiflora	For control. 2	1
	Treatments should be made prior to leaf deterioration by leaf-feeding insects.	
Russian olive	For partial control. 2 to 5	1 to 2
	For control. This product is not registered in California for use on Russian olive.	
Sage, black	For control.	1
	Thorough coverage of foliage is necessary for best results.	
Sage, white	For partial control. 2 to 5	1 to 2
Sagebrush, California	For control.	1
	Thorough coverage of foliage is necessary for best results.	
Salmonberry	For control. 2	1
Salt cedar	For partial control. 2 to 5	1 to 2
Sassafras	For partial control. 2 to 5	1 to 2
Sourwood	For partial control. 2 to 5	1 to 2
Sumac:, Poison, Smooth and Winged	For partial control. 2 to 5	1 to 2
Sweetgum	For control. 2 to 3	1 to 1.5
Swordfern	For partial control. 2 to 5	1 to 2
Tallowtree, Chinese	For control.	1
	Thorough coverage of foliage is necessary for best results.	
Tan oak resprouts	For suppression or partial control.	1
	Apply to resprouts less than 3 to 6 feet tall. Best results are obtained with fall applications.	
Thimbleberry	For control. 2	1
Tobacco, tree	For partial control.	1 to 2
	Thorough coverage of foliage is necessary for best results	
Trumpetcreeper	For control. 2 to 3	1 to 1.5
Waxmyrtle, southern	For partial control. 2 to 5	1 to 2
Willow	For control. 3	1

NONCROP USES

NONCROP, RECREATIONAL AND PUBLIC AREAS

Additional surfactant may be used. If additional surfactant is to be used follow the manufacturers rates and recommendations for use of the surfactant. When applied as directed for "NONCROP USES", under conditions described, this product controls annual and perennial weeds and woody brush listed on this label growing in areas such as airports, apartment complexes, ditch banks, dry ditches, dry canals, fencerows, golf courses, highways, industrial plant sites, lumber yards, manufacturing sites, office complexes, parking areas, parks, petroleum tank farms and pumping installations, pipelines, power and telephone rights-of-way, railroads, roadsides (guardrails, shoulders), schools, storage areas, utility substations and warehouse areas.

For specific rates of application and instructions for control of various annual and perennial weeds and woody brush, see the "WEEDS CONTROLLED" section of this label.

Chemical mowing - Perennials

This product will suppress perennial grasses listed in this section to serve as a substitute for mowing. Use 8 fluid ounces of this product per acre when treating tall fescue, fine fescue, orchardgrass or quackgrass covers. Use 6 fluid ounces of this product per acre when treating Kentucky bluegrass. Apply treatments in sufficient volume of water to ensure thorough coverage.

Use only in areas where some temporary injury or discoloration of perennial grasses can be tolerated.

Chemical mowing - Annuals

For growth suppression of some annual grasses, such as annual ryegrass, wild barley and wild oats growing in coarse turf on roadsides or other industrial areas, apply 4 to 5 fluid ounces of this product in sufficient volume of water to ensure thorough coverage.

Applications should be made when annual grasses are actively growing and before the seedheads are in the boot stage of development. Treatments may cause injury to the desired grasses.

RAILROADS

Bare ground, Ballast and Shoulders, Crossings, and Spot treatment

This product may be used to maintain bare ground on railroad ballast and shoulders. This product may be used to improve line-of-sight at railroad crossings and reduce the need for mowing along rights-of-way, wayside structures, and other similar areas. This product may be tank mixed with other herbicides for ballast, shoulder, spot, bare ground, and crossing treatments UNLESS SPECIFICALLY PROHIBITED BY THE PRODUCT LABEL.

Brush control

This product may be used to control woody brush and trees on railroad rights-of-way. Apply 4 to 10 quarts of this product per acre as a broadcast spray, using boom-type or boom-less nozzles. Up to 80 gallons of spray solutions per acre may be used. Apply a 3/4 to 2 percent solution of this product when using high-volume spray-to-wet applications. Apply a 5 to 10 percent solution of this product when using low volume directed sprays for spot treatment. This product may be tank mixed with other products for enhanced control of woody brush and trees UNLESS SPECIFICALLY PROHIBITED BY THE PRODUCT LABEL.

Actively Growing Bermudagrass and Bermudagrass Release

This product may be used to control or partially control many annual and perennial weeds for effective release of actively growing Bermudagrass. Apply 1 to 3 pints of this product in up to 80 gallons of spray solutions per acre. Use the lower rate when treating annual weeds below 6 inches in height (or runner length). Use the higher rate as weeds increase in size or as they approach flower or seedhead formation. These rates will also provide partial control of the following perennial species:

Bahiagrass	Fescue, tall	Trumpet creeper	Vaseygrass
Bluestem, silver	Johnsongrass		

This product may be tank-mixed with Spyder (Sulfometuron methyl) If tank-mixed, use no more than 1 to 3 pints of this product with 1 to 2 ounces of Spyder per acre. Use the lower rates of each product to control annual weeds less than 6 inches in height (or runner length) that are listed in this label and the Spyder label. Use the higher rates as annual weeds increase in size and approach the flower or seedhead stages. These rates will also provide partial control of the following perennial weeds:

Bahiagrass	Dallisgrass	Fescue, tall	Trumpet creeper
Blackberry	Dewberry	Johnsongrass	Vaseygrass
Bluestem, silver	Dock, curly	Poorjoe	Vervain, blue
Broomsedge	Dogfennel	Raspberry	

Use only on well-established bermudagrass. Bermudagrass injury may result from the treatment, but regrowth will occur under moist conditions. Repeat applications in the same season are not recommended, since severe injury may occur.

ROADSIDES

Shoulder treatments

This product may be used on road shoulders.

Guardrails and other obstacles to mowing

This product may be used to control weeds growing under guardrails and around signposts and other objects along the roadside.



Spot treatment

This product may be used as a spot treatment to control unwanted vegetation growing along roadsides.

Tank mixtures

This product may be tank mixed with other herbicides for shoulder, guardrail, spot and bare ground treatments UNLESS SPECIFICALLY PROHIBITED BY THE PRODUCT LABEL.

Release of Bermudagrass or Bahiagrass

Dormant applications

This product may be used to control or partially control many winter annual weeds and tall fescue for effective release of dormant bermudagrass or bahiagrass. Treat only when turf is dormant and prior to spring greenup. This product may also be tank-mixed with Spyder (Sulfometuron methyl) for residual control. Tank mixtures of this product with Spyder may delay greenup.

For best results on winter annuals, treat when plants are in an early growth stage (below 6 inches in height) after most have germinated. For best results on tall fescue, treat when fescue is at or beyond the 4- to 6-leaf stage.

Apply 8 to 64 fluid ounces of this product per acre alone or in a tank mixture with 1/4 to 1 ounce per acre of Spyder. Use only in areas where bermudagrass or bahiagrass are desirable ground covers and where some temporary injury or discoloration can be tolerated. To avoid delays in greenup and minimize injury, add no more than 1 ounce of Spyder per acre on bermudagrass and no more than 0.5 ounce of Spyder per acre on bahiagrass and avoid treatments when these grasses are in semi-dormant condition.

Actively growing bermudagrass

This product may be used to control or partially control many annual and perennial weeds for effective release of actively growing bermudagrass. Apply 1 to 3 pints of this product per acre. Use the lower rate when treating annual weeds below 6 inches in height (or runner length). Use the higher rate as weeds increase in size or as they approach flower or seedhead formation. These rates will also provide partial control of the following perennial species:

Bahiagrass	Fescue, tall	Trumpet creeper	Vaseygrass
Bluestem, silver	Johnsongrass		

This product may be tank-mixed with Spyder (Sulfometuron methyl). If tank-mixed, use no more than 1 to 2 pints of this product with 1 to 2 ounces of Spyder per acre. Use the lower rates of each product to control annual weeds less than 6 inches in height (or runner length) that are listed in this label and the Spyder label. Use the higher rates as annual weeds increase in size and approach the flower or seedhead stages. These rates will also provide partial control of the following perennial weeds:

Bahiagrass	Dallisgrass	Fescue, tall	Trumpet creeper
Bluestem, silver	Dock, curly	Johnsongrass	Vaseygrass
Broomsedge	Dogfennel	Poorjoe	Vervain, blue

Use only on well-established bermudagrass. Bermudagrass injury may result from the treatment, but regrowth will occur under moist conditions. Repeat applications of the tank mix in the same season are not recommended, since severe injury may occur.

Actively growing bahiagrass

For suppression of vegetative growth and seedhead inhibition of bahiagrass for approximately 45 days, apply 6 fluid ounces of this product per acre. Apply 1 to 2 weeks after full greenup or after mowing to a uniform height of 3 to 4 inches. This application must be made prior to seedhead emergence.

For suppression up to 120 days, apply 4 fluid ounces of this product per acre, followed by an application of 2 to 4 fluid ounces per acre about 45 days later. Make no more than 2 applications per year.

TANK MIXTURES FOR NONCROP SITES AND FORESTRY SITE PREPARATIONS

Razor Herbicide plus Spyder (Sulfometuron methyl)


Use on noncrop sites including airports, industrial plants, lumberyards, petroleum tank farms, pumping stations, railroads, roadsides, storage areas or other similar sites where bare ground is desired.

This tank mixture may also be used as a site preparation treatment for sites to be planted to jack pine, loblolly pine, red pine, slash pine, Virginia pine and other conifer species. When applied as directed for "NONCROP USES" under the conditions described, this product plus Spyder provides control of annual weeds listed in the "WEEDS CONTROLLED" section of the label for this product and control or partial control of the perennial weeds listed below.

Apply 1 to 2 quarts of this product with 2 to 4 ounces of Spyder per acre as a broadcast spray to actively growing weeds.

This mixture may be applied by aerial equipment in site prep operations. When applied by air, use the recommended rates.

This product plus Spyder tank mixtures may not be applied by air in California.



For control of annual weeds, use the lower rates of these products.

For control of the listed perennial weeds, use the higher rates of both products. For partial control, use the lower rates.

Bahiagrass	Dock, curly	Johnsongrass**	Trumpet creeper*
Bermudagrass*	Dogfennel	Poorjoe**	Vaseygrass
Broomsedge	Fescue, tall	Quackgrass	Vervain, blue

*Suppression at the higher rates only.

**Control at the lower rates.

Read and carefully observe the cautionary statements and all other information appearing on the labels of all herbicides used.

TANK MIXTURES NONCROP SITES

When applied as a tank mixture, this product provides control of the emerged annual weeds and partial control to the emerged perennial weeds listed in this label. THIS PRODUCT MAY BE TANK MIXED WITH MOST NONCROP HERBICIDES UNLESS PROHIBITED BY THE SPECIFIC LABEL.

Use according to the most restrictive label directions for each product in the mixture.

FARMSTEAD WEED CONTROL

When applied as directed for "NONCROP USES", under conditions described, this product controls undesirable vegetation listed on this label around farmstead building foundations, along and in fences, shelterbelts and for general nonselective farmstead weed control.

For specific rates of application and instructions for control of various annual and perennial weeds, see the "WEEDS CONTROLLED" section of this label.

FARM DITCHES

This product will suppress perennial grasses along farm ditches. Apply this product at a rate of 6 to 8 fluid ounces per acre. Use 8 fluid ounces per acre when treating tall (coarse) fescue, fine fescue, orchardgrass or quackgrass covers. For best suppression of these species, add ammonium sulfate at a rate of 1.7 pounds per 10 gallons of spray solution. Use 6 fluid ounces per acre without ammonium sulfate when treating Kentucky bluegrass.

Apply treatments to actively growing perennial grass covers. Recommended for best spray distribution and coverage, use flat fan nozzles. Additional surfactant may be used. If additional surfactant is to be used follow the manufacturers rates and recommendations for use of the surfactant.

Where broadleaf weed control or suppression is desired, tank mix this product with an appropriate, labeled broadleaf weed herbicide.

CONSERVATION RESERVE PROGRAM (CRP ACRES)

This product can be used to control undesirable vegetation when rotating out of CRP acres or to suppress competitive growth and seed production of undesirable vegetation in CRP acres.

For specific rates of application for various annual and perennial weeds, see the "WEEDS CONTROLLED" section of this label.

CRP applications may be made with wiper applicators or conventional spray equipment.

For selective applications with broadcast spray equipment, apply 12 to 16 ounces per acre of this product in early spring before desirable CRP grasses, such as crested and tall wheatgrass, break dormancy and initiate green growth. Late fall applications can be made after desirable perennial grasses have reached dormancy. Some stunting of CRP perennial grasses will occur if applications are made when plants are not dormant.

DORMANT RANGELAND

This product will control or suppress weeds in dormant rangeland. Refer to the "WEEDS CONTROLLED SECTION" of this label.

Apply 8 to 16 ounces per acre of this product in the early spring when the weeds have greened up, but desirable grasses, such as crested and tall wheatgrass are still truly dormant.

Slight discoloration of the desirable grasses may occur, but they will regreen and regrow under moist soil conditions as effects of this product wear off.

Do not use additional surfactant or ammonium sulfate when spraying dormant rangeland grasses with this product.

HABITAT MANAGEMENT

This product is recommended for the restoration and/or maintenance of native habitats and in wildlife management areas. Apply as recommended in the "NONCROP USES" section of this label.

Habitat Restoration and Maintenance - When applied as directed, exotic and other undesirable vegetation may be controlled in habitat management areas. Applications can be made to allow recovery of native plant species, prior to planting desirable native species, and for similar broad spectrum vegetation control requirements in habitat management areas. Spot treatments can be made to selectively remove unwanted plants for habitat maintenance and enhancement. For spot treatments, care should be exercised to keep spray off desirable plants.

Wildlife Food Plots -This product may be used as a site preparation treatment prior to planting wildlife food plots. Any wildlife food species may be planted after applying this product, or native species may be allowed to repopulate the area. If tillage is needed to prepare a seedbed, wait 7 days after applying this product before tilling.

ORNAMENTALS, NURSERIES (PLANTS AND TREES) AND CHRISTMAS TREES

THIS PRODUCT IS NOT RECOMMENDED FOR USE AS AN OVER-THE-TOP BROADCAST SPRAY IN ORNAMENTALS AND CHRISTMAS TREES.

Note: Desirable plants may be protected from the spray solution by using shields or coverings made of cardboard or other impermeable material.

When applied as instructed for the conditions described for "NONCROP USES", this product controls undesirable vegetation listed on this label prior to planting, within and around greenhouses and shadehouses, and as a postdirected spray around established ornamentals and Christmas trees. This product may also be used to trim and edge around trees, buildings, sidewalks, roads, potted plants and other objects in a nursery setting.

For specific rates of application and instructions for control of various annual and perennial weeds, see the "WEEDS CONTROLLED" section of this label.

Where repeat applications are necessary, do not exceed 10.6 quarts of this product per acre per year.

Site Preparation - Following preplant applications of this product, any ornamental, nursery species or Christmas tree species may be planted. Precautions should be taken to protect nontarget plants during site preparation applications.

Greenhouse/Shadehouse Use - This product may be used to control weeds listed on this label which are growing in greenhouses. Desirable vegetation must not be present during application and air circulation fans must be turned off.

Postdirected Spray - Use as a postdirected spray around established woody ornamental species, nursery species or Christmas trees such as those listed below. Care must be exercised to avoid contact of spray, drift or mist with foliage or green bark of established ornamental species.

Arborvitea	Euonymus	Lilac	Pine
Azalea	Fir	Magnolia	Spruce
Boxwood	Hollies	Maple	
Crabapple	Jobba	Oak, privet	

SILVICULTURAL SITES AND RIGHTS-OF-WAY

NOTE: NOT RECOMMENDED FOR USE AS AN OVER-THE-TOP BROADCAST SPRAY IN SILVICULTURAL NURSERIES

When applied as directed for "NONCROP USES" under conditions described, this product controls undesirable vegetation listed on this label.

For specific rates of application and instructions for control of various brush, annual and perennial weeds, see the "WEEDS CONTROLLED" section of this label.

Do not exceed 10.6 quarts of this product per acre per year.

Aerial Application - This product may be applied using aerial spray equipment for silvicultural site preparation, and rights-of-way treatments. See the "APPLICATION EQUIPMENT and TECHNIQUES" part of the "MIXING, ADDITIVES and APPLICATION INSTRUCTIONS" section of this label for information on how to apply this product by air.

DO NOT APPLY THIS PRODUCT BY AIR TO RIGHTS-OF-WAY SITES IN THE STATE OF CALIFORNIA.

SITE PREPARATION

Following preplant applications of this product, any silvicultural species may be planted.

POSTDIRECTED SPRAY

In established silvicultural sites, use as a spray on the foliage of undesirable vegetation. Care must be exercised to avoid contact of spray, drift or mist with foliage or green bark of desirable species.

CUT STUMP TREATMENTS

Woody vegetation may be controlled by treating freshly cut stumps of trees and resprouts with this product. Apply this product using suitable equipment to ensure coverage of the entire cambium. Cut vegetation close to the soil surface. Apply a 50 to 100 percent solution of this product to the freshly-cut surface immediately after cutting. Delays in application may result in reduced performance. For best results, applications should be made during periods of active growth and full leaf expansion.

When used according to directions for cut stump application, this product will CONTROL, PARTIALLY CONTROL or SUPPRESS many types of woody brush and tree species, some of which are listed below:

Alder	Oak	Salt cedar	Tan Oak
Eucalyptus	Reed, giant	Sweetgum	Willow
Madrone			

INJECTION AND FRILL APPLICATIONS

Woody vegetation may be controlled by injection or frill application of this product. Apply this product using suitable equipment which must penetrate into the living tissue. Apply the equivalent of 1 ml of this product per each 2 to 3 inches of trunk diameter (DBH). This is best achieved by applying a 50 to 100 percent concentration of this material either to a continuous frill around the tree or as cuts evenly spaced around the tree below all branches. As tree diameter increases in size, better results are achieved by applying diluted material to a continuous frill or more closely spaced cuttings. Avoid application techniques that allow runoff to occur from frill or cut areas in species that exude sap freely after frills or cutting. In species such as this, make frill or cut at an oblique angle so as to produce a cupping effect and use undiluted material. For best results, application should be made during periods of active growth and after full leaf expansion.

This treatment WILL CONTROL the following woody species:

Oak	Poplar	Sweetgum	Sycamore
-----	--------	----------	----------

This treatment WILL SUPPRESS the following woody species:

Blackgum	Dogwood	Hickory	Maple, red
----------	---------	---------	------------

TURFGRASSES AND GRASSES FOR SEED PRODUCTION

PREPLANT AND RENOVATION

When applied as directed for "NONCROP USES", under conditions described, this product controls most existing vegetation prior to the planting or renovation of either turfgrasses or grass seed production areas.

For specific rates of application and instructions for control of various annual and perennial weeds, and woody brush and trees, see the "WEEDS CONTROLLED" section of this label.

For maximum control of existing vegetation, delay planting to determine if any regrowth from escaped underground plant parts occurs. Where repeat treatments are necessary, sufficient regrowth must be attained prior to application. For warm-season grasses, such as bermudagrass, summer or fall applications provide best control.

DO NOT DISTURB SOIL OR UNDERGROUND PLANT PARTS BEFORE TREATMENT. Tillage or renovation techniques such as vertical mowing, coring or slicing should be delayed for 7 days after application to allow proper translocation into underground plant parts.

TURFGRASSES

Where existing vegetation is growing in a field or unmowed situation, apply this product to actively growing weeds at the stages of growth listed in the "WEEDS CONTROLLED" section of this label.

Where existing vegetation is growing under mowed turfgrass management in such sites as apartment complexes, residential areas and sod farms, apply this product after omitting at least one regular mowing to allow sufficient growth for good interception of the spray.

Desirable turfgrasses may be planted following the above procedures.

GRASSES FOR SEED PRODUCTION

Apply this product to actively growing weeds at the stages of growth recommended in the "WEEDS CONTROLLED" section of this label prior to planting or renovation of turf or forage grass areas grown for seed production.

DO NOT feed or graze treated areas within 8 weeks after application.

ANNUAL WEED CONTROL IN DORMANT BERMUDAGRASS AND BAHIAGRASS TURF

When applied as directed for "NONCROP USES" under the conditions described, this product will provide control or suppression of many winter annual weeds and tall fescue for effective release of dormant bermudagrass and bahiagrass turf. Refer to the rate table for Riverdale Razor alone under the "RELEASE OF BERMUDAGRASS and BAHIAGRASS" section of this label for recommended rates and volumes on the species to be suppressed or controlled. Treat only when turf is dormant and prior to spring greenup. Spot treatments or broadcast applications of this product in excess of 16 fluid ounces per acre may result in injury or delayed greenup in highly maintained turfgrass areas; i.e., golf courses, lawns, etc. DO NOT APPLY TANK MIXTURES of this product plus Spyder (Sulfometuron methyl) in highly maintained turfgrass areas.

RELEASE OF BERMUDAGRASS OR BAHIAGRASS

NOTE: Use only in areas where bermudagrass or bahiagrass are desirable ground covers and where some temporary injury or discoloration can be tolerated. Use tank mixtures of this product plus Spyder (Sulfometuron methyl) only on railroads, highways, utility plant sites, or other right-of-way areas.

When applied as directed for "NONCROP USES" under the conditions described, this product will provide control or suppression of many winter annual weeds and tall fescue for effective release of dormant bermudagrass or bahiagrass. This product may be tank-mixed with Spyder as recommended for residual control. Make applications to dormant bermudagrass or bahiagrass. Tank mixtures of this product plus Spyder may delay greenup. To avoid delays in greenup and minimize injury, do not add more than 1 ounce per acre of Spyder on bermudagrass or more than 0.5 ounce per acre on bahiagrass, or treat when these grasses are in a semi-dormant condition.

For best results on winter annuals, treat when plants are in an early growth stage (below 6 inches in height) after most have germinated. For best results on tall fescue, treat when fescue is in or beyond the 4 to 6-leaf stage.

WEEDS CONTROLLED

Rate recommendations for control or suppression of winter annuals and tall fescue are listed below:

Apply the recommended rates of this product alone or as a tank mixture in sufficient water to ensure thorough coverage.

For the best recommendation for the mixture of weeds within your geographic area, contact your Nufarm sales representative.

WEEDS CONTROLLED OR SUPPRESSED WITH RAZOR ALONE*						
WEED SPECIES	RIVERDALE RAZOR (Fluid Ounces/Acre)					
	8	12	16	24	32	64
Barley, little	S	C	C	C	C	C
Bedstraw, catchweed	S	C	C	C	C	C
Bluegrass, annual	S	C	C	C	C	C
Chervil	S	C	C	C	C	C
Chickweed, common	S	C	C	C	C	C
Clover, crimson	•	S	S	C	C	C
Clover, large hop	•	S	S	C	C	C
Fescue, tall	•	•	•	•	S	S
Geranium, Carolina	•	•	S	S	C	C
Henbit	•	S	C	C	C	C
Ryegrass, Italian	•	•	S	C	C	C
Speedwell, corn	S	C	C	C	C	C
Vetch, common	•	•	S	C	C	C

NOTE: C = Control
S = Suppression

*These rates apply only to sites where an established competitive turf is present.

WEEDS CONTROLLED OR SUPPRESSED WITH RIVERDALE RAZOR PLUS SPYDER (SULFOMETURON METHYL)* WEED SPECIES								
WEED SPECIES	RIVERDALE RAZOR (Fluid Ounces/Acre)	8	12	12	16	16	12	16
	+	+	+	+	+	+	+	+
	Spyder (Ounces/Acre)	1/4	1/4	1/2	1/4	1/2	1	1
Barley, little		C	C	C	C	C	C	C
Bedstraw, catchweed		C	C	C	C	C	C	C
Bluegrass, annual		S	C	C	C	C	C	C
Chervil		C	C	C	C	C	C	C
Chickweed, common		S	C	C	C	C	C	C
Clover, crimson		S	S	S	S	C	C	C
Clover, large hop		•	•	S	S	C	C	C
Fescue, tall		•	•	•	•	•	S	S
Geranium, Carolina		•	S	S	C	C	C	C
Henbit		•	S	C	C	C	C	C
Ryegrass, Italian		•	S	S	C	C	C	C
Speedwell, corn		S	C	C	C	C	C	C
Vetch, common		C	C	C	C	C	C	C

NOTE: C = Control
S = Suppression

*These rates apply only to sites where an established competitive turf is present.



RELEASE OF ACTIVELY GROWING BERMUDAGRASS

When applied as directed, this product will aid in the release of bermudagrass by providing control of annual species listed in the "WEEDS CONTROLLED" section of this and the Spyder (Sulfometuron methyl) label and suppression or partial control of certain perennial weeds.

For control or suppression of those annual species listed on this label, use 1 to 3 pints of this product as a broadcast spray in sufficient water to ensure thorough coverage. Use the lower rate when treating annual weeds below 6 inches in height (or length of runner in annual vines). Use the higher rate as weeds increase in size or as they approach flower or seedhead formation.

Use the higher rate of this product for partial control of the following perennial species. Use the lower rates for suppression of growth. For best results, see the "WEEDS CONTROLLED" section of this label for proper stage of growth.

Bahiagrass	Fescue, tall	Trumpetcreeper**	Vaseygrass
Bluestem, silver	Johnsongrass*		

*Control at the higher rates.

**Suppression at higher rates only.

This product may be tank-mixed with Spyder. If tank-mixed, use no more than 1 to 2 pints per acre of this product with 1 to 2 ounces of Spyder per acre.

Use the lower rates of both mixtures to control annual weeds below 6 inches in height (or runner length in annual vines) that are listed in the "WEEDS CONTROLLED" section of this booklet and the Spyder label. Use the higher rates as annual weeds increase in size and approach the flower or seedhead stages.

Use the higher rates of this product to provide partial control of the following perennial weeds. Use the lower rates for suppression of growth.

Bahiagrass	Dock, curly	Johnsongrass*	Vaseygrass
Bluestem, silver	Dogfennel	Poorjoe*	Vervain, blue
Broomsedge	Fescue, tall	Trumpetcreeper**	

*Suppression at higher rates only.

**Control at the higher rates.

COOL SEASON TURF GROWTH REGULATION

When applied as directed, this product will suppress growth and seedhead development of listed turf species in noncrop sites.

This product is recommended for management of coarse turf on roadside rights-of-way. Use in areas such as airports, apartment complexes, Christmas tree farms, ditch banks, dry ditches, dry canals, fencerows, golf courses, highways, industrial sites, lumberyards, manufacturing sites, office complexes, ornamental nurseries, parks, parking areas, pipelines, petroleum tank farms and pumping installations, railroads, recreational pipeline areas, residential areas, rights-of-way, roadsides (including guardrails and shoulders), sod or turf seed farms, schools, storage areas, substations and warehouse areas. Do not use on high-quality turf or other areas where some turf color changes cannot be tolerated. Slight turf discoloration may occur but turf will regreen and regrow under moist conditions as effects of this product wear off.

Apply 4 to 6 fluid ounces of this product per acre alone or in a recommended tank mixture.

Apply the recommended rates of this product alone or as a tank mixture in sufficient water to ensure thorough coverage.

This product can be used for growth and seedhead suppression of:

TALL FESCUE/SMOOTH BROME

For best results, apply this product in a recommended tank mixture to actively growing turfgrasses after greenup in the spring of the year. For suppression of seedheads, applications must be made before boot-to-seedhead stage of development. Applications made from seedhead emergence until maturity may result in turf discoloration or injury.

After mowing or removal of seedheads, this product in a recommended tank mixture may also be used to suppress the growth of certain turfgrasses. Allow turf to recover from stress caused by heat, drought or mowing before making applications. Applications made to turf under stress may increase the potential for discoloration or injury.

ANNUAL GRASSES


For growth suppression of some annual grasses such as annual ryegrass, wild barley and wild oats, apply 3 to 4 fluid ounces of this product per acre. Applications should be made when annual grasses are actively growing and before the seedheads are in the boot stage of development. Treatments made after seedhead emergence may cause injury to the desired grasses.

TANK MIXTURES

For the following tank mixtures, consult each product label for weeds controlled and the correct stage of application. Do not treat turf under stress.

Tank mixtures plus 2,4-D Amine

For additional weed control benefits, up to 1 pound a.i. per acre of 2,4-D amine may be added to the following tank mixtures. Consult the label for 2,4-D amine for weeds controlled.



TALL FESCUE**This product plus Corsair™ (Clorsulfuron)**

For suppression of tall fescue growth and seedheads, and control or partial control of some annual weeds, apply this tank mixture after greenup and prior to boot-to-seedhead stage of development. Use up to 0.5 ounce of clorsulfuron per acre.

This tank mixture can also be applied after mowing or removal of tall fescue seedheads for turf growth suppression. Make only one of the above applications per growing season.

This product plus Spyder (Sulfometuron methyl)

For suppression of tall fescue growth and seedheads, and control or partial control of some annual weeds, apply this tank mixture after greenup and prior to boot-to-seedhead stage of development. Use up to 0.25 ounce of sulfometuron methyl per acre.

This product plus Manor™ (Metsulfuron methyl)

This tank mixture can be applied after mowing or removal of tall fescue seedheads for turf growth suppression and control or partial control of some annual weeds. Use up to 1/3 ounce of metsulfuron methyl per acre.

SMOOTH BROME**This product plus Spyder (Sulfometuron methyl)**

For suppression of smooth brome growth and seedheads and control or partial control of some annual weeds, apply this tank mixture after greenup and prior to boot-to-seedhead stage of development. Use up to 0.25 ounce of sulfometuron methyl per acre.

STORAGE AND DISPOSAL

Do not contaminate water, foodstuffs, feed or seed by storage or disposal.

PESTICIDE STORAGE: STORE ABOVE 10°F(-12°C) TO KEEP PRODUCTS FROM CRYSTALLIZING. Crystals will settle to the bottom. If allowed to crystallize, place in a warm room 68°F(20°C) for several days to redissolve and shake, roll or agitate to mix well before using.

PESTICIDE DISPOSAL: Wastes resulting from the use of this product that cannot be used or chemically reprocessed should be disposed of in a landfill approved for pesticide disposal or in accordance with applicable Federal, state or local procedures. Emptied container retains vapor and product residue. Observe all label safeguards until container is destroyed.

CONTAINER DISPOSAL: Do not reuse container. Triple rinse container. Then puncture and dispose of in a sanitary landfill, or by incineration, or, if allowed by state and local authorities, by burning. If burned, stay out of smoke.

CONTAINER DISPOSAL FOR REFILLABLE CONTAINERS: Close all openings which have been opened during use and replace all caps. Contact Nufarm Customer Service at 1-800-345-3330, to arrange for return of the empty refillable container.

WARRANTY

Seller warrants that the product conforms to its chemical description and is reasonably fit for the purpose stated on the label when used in accordance with directions under normal conditions of use, but neither this warranty nor any other warranty of merchantability or fitness for a particular purpose, expressed or implied, extends to the use of this product contrary to label instructions, or under abnormal conditions or under conditions not reasonably foreseeable to seller, and buyer and the limit of liability of any such use. The exclusive remedy of user or buyer and the limit of liability of Nufarm Americas, Inc. is the purchase price paid for the quantity of product involved.

(RV111804)

Razor and Riverdale are registered trademarks and Corsair, Diablo, Manor and Spyder are trademarks of Nufarm Americas Inc.
Weedar and Weedone are registered trademarks of Nufarm, Inc.

Frontier, Guardsman, Marksman, Prowl, Pursuit, Pursuit Plus, Scepter, Solicam and Squadron are registered trademarks of BASF, Inc.

DEF, Sencor and Turbo are registered trademarks of Bayer AG.

Folex and Prep are registered trademarks of Bayer Crop Sciences.

Broadstrike, Surflan, Surpass and Topnotch are registered trademarks and First Rate, Frontrow, Pendimax and Python are trademarks of Dow AgriScience LLC.

Bladex, Canopy, Extrazine, Gemini, Karmex, Krovar, Lexone, Lorox, Lorox Plus, and Preview are registered trademarks of E.I. duPont de Nemours and Company.

Command is a registered trademark of FMC Corporation.

Linex is a registered trademark of Griffin LLC.

Bullet, Harness, Lasso, Lariat, Micro-Tech, Partner are registered trademarks of Monsanto Company.

Princep and Caliber are trademarks of Novartis Corporation.

Goal is a registered trademark of Rohm and Haas Company.

Bicep, Dual, Caliber and Princep are registered trademarks of Syngenta Crop Protection.

Fusion is a registered trademark of ZENECA Company.

This product is protected by U.S. Patent No. 4,405,531. Other patents pending. No license granted under any non-U.S. patent(s).

RIVERDALE® RAZOR® HERBICIDE

THE COMPLETE BROAD SPECTRUM POSTEMERGENCE PROFESSIONAL HERBICIDE FOR INDUSTRIAL, FORESTRY, TURF, VEGETATION MANAGEMENT AND ORNAMENTAL WEED CONTROL

ACTIVE INGREDIENT:	Glyphosate, N-(phosphonomethyl) glycine, in the form of its isopropylamine salt*.....	41.0%
OTHER INGREDIENTS:	59.0%
	TOTAL.....	100.0%

* Contains 480 grams per litre or 4 pounds per U.S. gallon of the active ingredient glyphosate, in the form of its isopropylamine salt. Equivalent to 356 grams per litre or 3 pounds per U.S. gallon of the acid, glyphosate.

KEEP OUT OF REACH OF CHILDREN
CAUTION - CAUCION

Si usted no entiende la etiqueta, busque a alguien para que se la explique a usted en detalle.
(If you do not understand the label, find someone to explain it to you in detail.)

SEE ATTACHED BOOKLET FOR ADDITIONAL PRECAUTIONARY STATEMENTS

For Chemical Spill, Leak, Fire, or Exposure, Call CHEMTREC (800) 424-9300. For Medical Emergencies Only, Call (877) 325-1840.

PRECAUTIONARY STATEMENTS
HAZARDS TO HUMANS AND DOMESTIC ANIMALS
CAUTION - CAUCION

Causes moderate eye irritation. Harmful if swallowed or inhaled. Do not get in eyes or on clothing. Avoid breathing vapor or spray mist.

FIRST AID	
IF IN EYES	<ul style="list-style-type: none">• Hold eye open and rinse slowly and gently with water for 15 to 20 minutes.• Remove contact lenses, if present, after the first 5 minutes, then continue rinsing eye.• Call a poison control center or doctor for treatment advice.
IF SWALLOWED	<ul style="list-style-type: none">• Call a poison control center or doctor immediately for treatment advice.• Have person sip a glass of water if able to swallow.• Do not induce vomiting unless told to do so by a poison control center or doctor. Do not give anything by mouth to an unconscious person.
IF INHALED	<ul style="list-style-type: none">• Move person to fresh air.• If person is not breathing, call 911 or an ambulance then give artificial respiration, preferably mouth-to-mouth if possible.• Call a poison control center or doctor for further treatment advice.
HOT LINE NUMBER	
Have the product container or label with you when calling a poison control center or doctor, or going for treatment. You may also contact 1-877-325-1840 for emergency medical treatment information.	

ENVIRONMENTAL HAZARDS

Do not apply directly to water, to areas where surface water is present or to intertidal areas below the mean high water mark. Do not contaminate water when disposing of equipment washwaters.

DIRECTIONS FOR USE

It is a violation of Federal law to use this product in a manner inconsistent with its labeling. READ ENTIRE LABEL BEFORE USING THIS PRODUCT. Do not apply this product in a way that will contact workers or other persons, either directly or through drift. Only protected handlers may be in the area during application. For any requirements specific to your State or Tribe, consult the agency responsible for pesticide regulations.

STORAGE AND DISPOSAL

Do not contaminate water, foodstuffs, feed or seed by storage or disposal.

PESTICIDE STORAGE: STORE ABOVE 10°F (-12°C) TO KEEP PRODUCTS FROM CRYSTALLIZING. Crystals will settle to the bottom. If allowed to crystallize, place in a warm room 68°F (20°C) for several days to redissolve and shake, roll or agitate to mix well before using.

PESTICIDE DISPOSAL: Wastes resulting from the use of this product that cannot be used or chemically reprocessed should be disposed of in a landfill approved for pesticide disposal or in accordance with applicable Federal, state or local procedures. Emptied container retains vapor and product residue. Observe all label safeguards until container is destroyed.

CONTAINER DISPOSAL: Do not reuse container. Triple rinse container. Then puncture and dispose of in a sanitary landfill, or by incineration, or, if allowed by state and local authorities, by burning. If burned, stay out of smoke.

CONTAINER DISPOSAL FOR REFILLABLE CONTAINERS: Close all openings which have been opened during use and replace all caps. Contact Nufarm Customer Service at 1-800-345-3330, to arrange for return of the empty refillable container.

MANUFACTURED BY
NUFARM AMERICAS INC.
BURR RIDGE, IL 60527-0866

EPA REG. NO. 228-366
EPA EST. NO. 228-IL-1

PULL HERE TO OPEN
PRESS TO RESEAL

331015

RV030905

3667-RIV